
LARGE LANGUAGE MODELS-ENABLED DIGITAL TWINS FOR PRECISION MEDICINE IN RARE GYNECOLOGICAL TUMORS

A PREPRINT

Jacqueline Lammert^{*1,2,3,4}, Nicole Pfarr^{2,5}, Leonid Kuligin⁶, Sonja Mathes^{7,8}, Tobias Dreyer^{1,3}, Luise Modersohn⁹, Patrick Metzger¹⁰, Dyke Ferber^{11,12}, Jakob Nikolas Kather^{11,12,13}, Daniel Truhn¹⁴, Lisa Christine Adams¹⁵, Keno Kyrill Bressen¹⁶, Sebastian Lange^{2,17}, Kristina Schwamborn^{2,5}, Martin Boeker⁹, Marion Kiechle¹, Ulrich A. Schatz^{1,2}, Holger Bronger^{1,3}, and Maximilian Tschochohei⁶

¹Department of Gynecology and Center for Hereditary Breast and Ovarian Cancer, Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

²Center for Personalized Medicine (ZPM), Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

³German Cancer Consortium (DKTK), partner site Munich, a partnership between DKFZ and TUM University Hospital, Germany

⁴European Reference Network for RARE CANcers (EURACAN) Initiative, partner site Munich, Germany

⁵Institute of Pathology, Technical University of Munich (TUM), School of Medicine and Health, Munich, Germany

⁶Google Cloud, Munich, Germany

⁷Department of Dermatology and Allergy Biederstein, Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

⁸Institute for History, Theory and Ethics of Medicine, University of Mainz Medical Center, Mainz, Germany

⁹Institute of Artificial Intelligence in Medicine and Healthcare, Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

¹⁰Institute of Medical Bioinformatics and Systems Medicine, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

¹¹Else Kroener Fresenius Center for Digital Health, Technical University Dresden, Dresden, Germany

¹²National Center for Tumor Diseases, Heidelberg University Hospital, Heidelberg, Germany; Department of Medical Oncology, Heidelberg University Hospital, Heidelberg, Germany

¹³Department of Medicine I, University Hospital Dresden, Dresden, Germany

¹⁴Department of Diagnostic and Interventional Radiology, University Hospital RWTH Aachen, Aachen, Germany

¹⁵Department of Diagnostic and Interventional Radiology, Technical University of Munich, School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

¹⁶Department of Cardiovascular Radiology and Nuclear Medicine, Technical University of Munich, School of Medicine and Health, German Heart Center, TUM University Hospital, Munich, Germany

¹⁷Department of Medicine II, Technical University of Munich (TUM), School of Medicine and Health, Klinikum rechts der Isar, TUM University Hospital, Munich, Germany

September 4, 2024

ABSTRACT

Rare gynecological tumors (RGTs) present major clinical challenges due to their low incidence and heterogeneity. The lack of clear guidelines leads to suboptimal management and poor prognosis. Molecular tumor boards accelerate access to effective therapies by tailoring treatment based on biomarkers, beyond cancer type. Unstructured data that requires manual curation hinders efficient use of biomarker profiling for therapy matching. This study explores the use of large language models (LLMs) to construct digital twins for precision medicine in RGTs.

Our proof-of-concept digital twin system integrates clinical and biomarker data from institutional and published cases (n=21) and literature-derived data (n=655 publications with n=404,265 patients) to

*Corresponding author: Jacqueline Lammert, MD, BSc Mail: jacqueline.lammert@tum.de

create tailored treatment plans for metastatic uterine carcinosarcoma, identifying options potentially missed by traditional, single-source analysis. LLM-enabled digital twins efficiently model individual patient trajectories. Shifting to a biology-based rather than organ-based tumor definition enables personalized care that could advance RGT management and thus enhance patient outcomes.

Keywords Large Language Models (LLMs) · Digital Twins · Precision Oncology · Rare Gynecological Tumors (RGTs)

1 Introduction

Rare Gynecological Tumors (RGTs), comprising over 30 distinct histological subtypes, such as sex cord stromal tumors, and uterine or ovarian carcinosarcomas, account for more than 50% of gynecologic malignancies, presenting a major clinical challenge.¹ With an incidence rate below six per 100,000 individuals, RGTs are difficult to study through large-scale randomized trials, leading to reliance on less standardized treatment approaches such as retrospective studies, case reports, and expert opinions. This lack of robust clinical guidelines has contributed to persistently poor prognosis for patients with RGTs.²

Technological advancements in cancer diagnostics have enabled the identification of biomarkers as therapeutic targets. Biomarker-guided treatments promise to accelerate the development of precision therapeutics across tumor types, reducing the relevance of organ-based classification.³ The prevailing organ-centric approach to clinical trial design hinders the development of effective treatments for rare cancers with shared biomarkers.⁴ This obstacle extends beyond rare cancers: The premature dismissal of olaparib in ovarian cancer and the seven to ten year delay in addressing Programmed Cell Death Ligand 1 (PD-L1) expressing breast and gynecological cancers with PD-L1 inhibition illustrate the need for biomarker-driven stratification for cancer treatment.^{5,6}

Molecular tumor boards (MTBs) are essential for interpreting biomarker profile results and matching cancer patients with appropriate therapies. This includes identifying suitable investigational drugs.⁷ The manual interpretation of multiple, co-occurring molecular alterations requires an in-depth understanding of their functional implications and correlations with treatment sensitivity or resistance. The rapid growth of biomedical literature and the fragmented nature of data sources make manual curation a bottleneck in efficiently translating genomic data into actionable treatment strategies.⁷

The data produced by MTBs is often stored in unstructured formats within electronic health records (EHRs) or other repositories, hindering their reusability for similar patients. Evaluating the effectiveness of MTB-guided treatments requires extracting follow-up data from EHRs. Unstructured text within EHRs, coupled with the lack of interoperability across healthcare institutions – particularly when MTB patients receive treatment at external facilities – renders the process labor-intensive, error-prone, and time-consuming.⁸ Consequently, applying MTB insights to future patients is hindered.

Advances in data capture and analysis, alongside decreasing costs in genome sequencing, are paving the way for innovative tools to manage rare or refractory cancers more effectively.⁹ Digital twin technology constructs virtual representations of physical entities with dynamic, bidirectional interfaces.¹⁰ Initially applied in industrial engineering, digital twins can also represent the human body in healthcare. By modeling physiological processes and predicting biomarker-specific responses to treatments, digital twins can address the challenges of patient variability and the limitations of traditional one-size-fits-all approaches.¹¹ In the case of RGT, the standard carboplatin and paclitaxel regimen, followed by chemotherapy monotherapies for subsequent lines, may not be the most effective approach.¹² Digital twins could help stratify RGT patients based on their unique biomarker profiles, enabling more tailored treatments and potentially improving outcomes, even in heavily pretreated cases.

Despite their potential, the adoption of digital twins in clinical practice is constrained by the challenges associated with integrating the diverse and complex data required for their development.¹³ Large language models (LLMs) offer potential to assist in this process by efficiently extracting and synthesizing relevant information from diverse sources.¹⁴

In this study, we demonstrate the application of an LLM-enabled workflow for constructing digital twins for patients with RGT, specifically metastatic uterine carcinosarcoma (UCS).

The research question was inspired by a real-world UCS case presented to a major German cancer center for evaluation of third-line treatment options. According to a consensus statement by Bogani et al., third-line monotherapy in UCS typically results in a median progression-free survival (PFS) of 1.8 months and a response rate of 5.5%, highlighting the urgent need for novel therapeutic strategies.¹² The patient presented with a proficient mismatch repair (pMMR) carcinosarcoma with intermediate Tumor Mutational Burden (TMB) and high PD-L1 expression. Although PD-L1 positivity is common in UCS^{15,16} and has been suggested as an independent prognostic factor,¹⁷ it has not been validated as a target for immunotherapy.¹²

Given the potential therapeutic importance of PD-L1, we investigated outcomes in similar patients. We identified cases with high PD-L1 expression, pMMR status and low to intermediate TMB from the institutional MTB database, including non-gynecological cancers, and to expand the pool of UCS cases, from the literature. The unstructured nature of EHR and academic publications posed challenges for immediate analysis. We utilized a local LLM to extract and structure data from EHRs and a cloud-based LLM for literature data. These datasets were integrated into a unified local database, forming the foundation of an RGT Digital Twin system. This system enabled the generation of virtual representations of individual patients, allowing for the simulation of personalized treatment strategies.

The RGT Digital Twin system facilitated the identification of additional therapeutic options, which were subsequently evaluated by MTB members. By integrating data from institutional sources (including non-gynecological cancers) and literature sources (to expand the pool of UCS cases), this approach provided novel insights that were not apparent from either data source alone. This integration has the potential to guide more effective treatment strategies for RGT patients and supports a shift towards a biology-based rather than organ-based definition of tumors. LLM technology enabled us to streamline the extraction, structuring, and analysis of EHR and web data, making it readily accessible for MTB evaluation. This is especially valuable in resource-limited settings like MTBs, where results can occasionally arrive too late to guide timely treatment decisions.¹⁸

2 Methods

2.1 Study setup

We employed an RGT Digital Twin system to create personalized treatment suggestions for UCS. A real-world patient case, along with molecular profiling data, was analyzed using the RGT Digital Twin system. The findings were then compared to analogous cases drawn from institutional and public databases. Treatment options were discussed at the MTB to inform individualized care decisions. Post-treatment outcomes were documented in the patient's EHR and updated for the individual RGT Digital Twin to improve future predictions. The RGT Digital Twin provided rationale for cost coverage requests and supported study inclusion decisions. Refer to Figure 1 for an overview of the study process.

2.2 Patient description and research question

The patient is a 77-year-old woman with metastatic UCS, initially diagnosed with FIGO IIIC2 UCS at age 66. Six years after surgery and adjuvant chemotherapy with carboplatin and paclitaxel, the patient experienced a recurrence in the cervical lymph nodes and pelvis. A cervical lymph node biopsy confirmed the recurrence, and the patient underwent the same chemotherapy regimen followed by MTB presentation in 2021 (see Supplementary Table 1 for detailed results). Genomic profiling was conducted using the TruSight Oncology 500 (TSO 500) and TruSight Tumor 170 (TST 170) panels. By analyzing a wide range of cancer-related genes, these panels facilitate the discovery of potential therapeutic targets. Molecular profiling revealed an intermediate TMB of 6.3 mutations/megabase, high PD-L1 expression (Combined Positive Score, CPS: 41), and a pMMR status. The patient's high PD-L1 expression prompted us to investigate the potential efficacy of PD-L1 inhibitors in metastatic tumors, regardless of primary tumor site or regional drug approval status. To this end, we searched our institutional MTB database for analogous cases, not restricted to gynecological cancers, and expanded our cohort with additional UCS cases identified in the literature.

2.3 Data collection

EHR data obtained from Technical University of Munich (TUM) University Hospital was the primary data source. Data downloaded from web-based repositories through institutional access extended the dataset. These repositories included *PubMed*, *ClinicalTrials.gov*, and clinical practice guidelines from the National Comprehensive Cancer Network (NCCN) and the German Cancer Society (DKG), which ensured adherence to up-to-date clinical standards.

A two-stage approach was employed for extracting structured, actionable data from source files. First, a locally deployed LLM system extracted relevant data from institutional EHRs. Second, a cloud-based LLM system processed documents from web-based repositories. Afterwards, the extracted structured dataset was made available to clinicians and the locally deployed LLM for further analysis. This process is shown in Figure 2.

2.4 Identification of analogous institutional cases

Analysis of the institutional MTB database at the TUM University Hospital identified cases analogous to the presented UCS patient. The analysis included patients discussed at the MTB between September 2017 and July 2024. Eligibility

criteria for screening included high PD-L1 expression ($CPS \geq 40$) and availability of MMR and TMB status. To avoid bias, we excluded patients with high TMB and deficient Mismatch Repair, which is known to be responsive to ICI and approved as Food and Drug Administration targets for ICI therapy. Similarity to the UCS case was determined based on medical discipline (gynecological oncology), or histopathological features independent of gender or origin (carcinosarcoma or sarcomatoid carcinoma morphology). Given the shared molecular and genomic characteristics between UCS and high-grade serous ovarian and endometrial carcinomas, gynecological cancers were chosen as a criterion for similarity.¹² Due to the unclear clinical utility of distinguishing carcinosarcoma from sarcomatoid carcinoma, the institutional MTB members combined them under the category morphology. Patients were included in the final analysis if they met all of the following conditions: $CPS \geq 40$, pMMR status, $TMB < 15$ mutations/megabase, and conformance to at least one of the predefined similarity parameters.

2.5 Institutional patient data extraction pipeline

EHR of selected patients were processed in a secure hospital environment. Documents varied in format, from (handwritten) medical notes to obituaries. Ten attributes were extracted from documents for each patient to form the RGT Digital Twin. Supplementary Table 2 shows the full list of attributes. Optical Character Recognition (OCR) was performed using Tesseract.¹⁹ Raw text was processed with a locally deployed version of pre-trained LLM *gemma-2-27b-it*, chosen for its ability to run locally while maintaining strong performance on medical texts.²⁰ This privacy-preserving architecture ensured that patient data would not leave the local clinic environment. In-context learning was used to adapt the LLM to the task at hand. With this method, LLMs receive extensive instructions in their prompt, e.g., in the form of example input and output. This improves their recall and precision.²¹ The study was approved by the local ethics committee of TUM (Reference No. 2023-486-S-SB).

2.6 Literature extraction pipeline

To extend the limited sample of analogous digital twins available in institutional EHR, a systematic literature search was conducted on *PubMed* using the terms ‘uterine carcinosarcoma’ and ‘endometrial carcinosarcoma’. Studies and case reports that included individual clinical follow-up data on patients with UCS treated with ICI were considered for inclusion in the analysis. Potential alternative treatment options and therapeutic targets were identified through a comprehensive review of national (DKG) and international (NCCN) oncological guidelines, *PubMed*-indexed publications, as well as the *ClinicalTrials.gov* database.

Data points were extracted in the structure shown in Supplementary Table 2. The sample size was captured as an additional data point. Additionally, the LLM was instructed to extract the main treatment recommendation based on the patient profile in the paper. General purpose LLM *Google-Gemini-1.5-Pro* was selected for this task due to its large context window, which enabled it to process all files in the sample without splitting them into smaller chunks.²² Since no institutional patient data was processed in this step, use of public cloud resources was permitted. The LLM was instructed to return results in the form of a JSON object. The LLM processed all documents sequentially, with each document processed in-context. Outputs were exported to a Pandas dataframe on the local machine in the secure hospital environment for convenient analysis by clinicians.

2.7 Construction of LLM-Enabled Digital Twin System

Next, extracted data points were stored in a database in the secure hospital environment that constituted the patient’s digital twin. Clinicians were able to model potential outcomes for their patients and determine suitable treatment strategies by reviewing treatment outcomes from patients with similar biomarkers and treatment history. Additionally, they were able to employ the local LLM to combine treatment strategies identified from web sources with the patient’s digital twin, creating personalized treatment recommendations. After selecting a treatment strategy, the database served as evidence for MTB evaluation, clinical trial matching, and creation of cost coverage requests with health insurance providers.

The developed pipeline is illustrated in Figure 2.

2.8 Analysis

Clinical characteristics, treatment regimens, duration of therapy, treatment responses, PFS, and overall survival (OS) were systematically collected from patient records and reports. Treatment response was captured from radiology reports and categorized as complete response (CR), partial response (PR), stable disease (SD), mixed response (MR), or progressive disease (PD). For additional treatment strategies, outcomes were summarized for each therapeutic approach.

Cases were sequentially numbered, starting with those retrieved from the institutional MTB database, followed by cases identified from the literature.

Formal statistical analysis to evaluate the accuracy of LLM data retrieval was performed by experts. Due to the large amount of data processed by the Digital Twin pipeline, we adopted human-in-the-loop reviews, an important aspect of machine learning.²³ To ensure that no information was missed during extraction, a sample-based review of LLM output was performed according to machine learning leading practice.²⁴ For institutional data, experts reviewed all attributes extracted from EHR by the LLM for correctness. For public research data, experts reviewed a random sample of attributes extracted from scientific studies for correctness. Additionally, all attributes that were used by the LLM to construct the literature-derived digital twins were manually reviewed. Afterwards, accuracy, precision, recall, and F1 scores of LLM extraction were calculated. Finally, all treatment recommendations generated by the LLM were manually reviewed and corrected by human experts. The data extraction review panel included two bioinformaticians and two gynecological oncologists with five and 16 years of clinical experience.

A panel of five MTB members, including three clinicians, one pathologist, and one biologist, along with a senior gynecological oncologist, evaluated the personalized treatment recommendations generated by the RGT Digital Twin system.

All statistical analyses were conducted using Pandas and SciPy libraries in Python (Version 3.10.12). The full code and documentation is available on GitHub.

3 Results

3.1 Patient cohort

A retrospective analysis of 1821 cases discussed at the institutional MTB between September 2017 and July 2024 was conducted. Among these, 132 cases exhibited high PD-L1 expression (CPS ≥ 40), encompassing 28 different tumor entities. The analysis was restricted to patients with TMB < 15 mutations/megabase and pMMR status with either gynecological cancers or carcinosarcoma/sarcomatoid carcinoma, resulting in a cohort of nine patients. Of these, seven patients received ICI therapy and were included in the study. The cohort comprised six females and one male aged 32 to 83 years at MTB presentation. Given that the similarity analysis focused on biomarker profiles and cancer morphology, the male patient's inclusion was appropriate. His sarcomatoid carcinoma aligned with the other inclusion criteria, regardless of his gender or cancer type.

3.2 Data extraction

89 EHR documents were extracted for the patient cohort (median: 11, range: 9-21). Documents had a median of two pages and 4,340 characters. Documents contained 70 data points for the selected attributes. Experts reviewed all extractions in the sample. The local LLM achieved accuracy of 0.76, precision of 0.96, recall of 0.78 and F1 of 0.86. The highest accuracy was achieved for 'diagnosis' and 'ICI treatment' (1.00). Low recall occurred in 'previous treatment' (0.29) and 'PFS' (0.14), mainly due to parsing errors in order and dates of previous treatments. See Table 2 for full results of the analysis.

Document analysis revealed that primary tumor sites included metastatic UCS (n=1), metastatic cervical cancer (n=4; three squamous cell carcinoma, one adenocarcinoma), metastatic uterine serous carcinoma (n=1), and metastatic, undifferentiated sarcomatoid carcinoma of the pancreas (n=1). Patients exhibited high PD-L1 expression with a median CPS of 75 (range: 40-95) and a median TMB of 5.5 (range: 0-11). Median follow-up duration was 48 months (range: 15-132 months). Detailed baseline clinical characteristics are presented in Supplementary Table 3.

The LLM-based systematic literature research yielded a dataset of 663 scientific documents. Files had a median of seven pages and 27,995 characters, with a maximum of 934,513 characters. The LLM extracted 7,956 attributes from scientific documents. Attribute extraction was reviewed with a random sample of n = 352 (Z = 1.96, N = 7,956, e = 0.05, P = 0.5). The cloud-based LLM achieved accuracy of 0.98, precision of 1.00, recall of 0.97, and F1 of 0.98. Lowest recall was observed in 'PFS' (0.77).

The LLM system identified 15 studies reporting ICI treatment in UCS, encompassing a total of 215 patients. While seven of the studies did not exclusively enroll UCS patients, four provided stratified outcomes for UCS cases. Phase II studies that provided stratified analysis for UCS patients showed objective response rates between zero and ten percent for ICI treatment. None of the seven studies allowed for individual patient-level data extraction to create digital twins (see details on these seven studies in Supplementary Table 4).

PD-L1 status was reported in 10 of the 215 literature-derived UCS cases treated with immunotherapy, with three cases exhibiting PD-L1 positivity. This limited sample size precluded stratified analysis. Notably, two of the PD-L1-positive UCS patients harbored dMMR and one had a high TMB, both of which are known to influence ICI treatment response.

Further analysis of the 15 identified studies yielded eight studies with individual patient follow-up data, comprising a total of 14 cases. The median age of these literature-derived patients was 63 years (range: 55-68 years).

3.3 Treatment response outcomes for 21 individual patients

In the institutional cohort, seven patients received ICI therapy: five with pembrolizumab monotherapy, one with pembrolizumab plus lenvatinib, and one with ipilimumab plus nivolumab. ICI therapy was initiated on average in the third line (range: 2-4). Median PFS was 6 months (range: 1-48). One patient remained disease-free after 45 months, two continued to respond, one received a subsequent treatment line, and three had died.

Treatments in the 14 cases of the literature-derived cohort consisted of pembrolizumab (n=4), pembrolizumab plus lenvatinib (n=7), pembrolizumab plus lenvatinib plus letrozole (n=1), PD-1/Cytotoxic T-lymphocyte associated protein 4 inhibitors (n=1), and avelumab plus axitinib (n=1). ICI was typically given in the third line (range: 2-5). Median PFS was 4 months (range: 0.9-15), and median OS was 9.9 months (range: 2.1-48). At data cut-off, six patients were alive, seven had died, and one had unknown status.

Table 2 provides a summary of ICI treatment response outcomes for all 21 cases.

3.4 RGT Digital Twins enable predictive modeling of individualized patient treatment strategies

To inform personalized treatment planning for the UCS patient (case 1), digital twins were created based on 21 evaluable patients. Treatment outcomes were predicted based on a database of additional 404,265 cases derived from scientific papers (n = 655). Potential treatment strategies were predicted for a patient with UCS with disease progression following third-line pembrolizumab monotherapy.

Supplementary Figure 1 presents treatment-relevant biomarkers after progression on standard-of-care combination treatment with carboplatin and paclitaxel identified by the digital twin system.

The digital twin system tailored treatment recommendations based on the patient's specific tumor characteristics, treatment history, and geographic location. Considering the patient's ongoing pembrolizumab therapy, the system suggested testing for Folate Receptor Alpha (FR α) to assess potential eligibility for an off-label treatment regimen currently under clinical investigation. This trial investigated the combination of mirvetuximab soravtansine and pembrolizumab in FR α -positive UCS, with eligibility criteria including pMMR status and prior pembrolizumab progression. However, the trial was no longer recruiting participants and was limited to the United States.²⁵ Therefore, the digital twin system suggested considering off-label use of this regimen for the patient. A previous evaluation (2021) identified HER2 amplification in the patient's tumor, a biomarker linked to high objective response rates to trastuzumab deruxtecan.²⁶ Due to the potential for HER2 status to evolve, the system recommended confirming this finding through a new biopsy.²⁷ Additionally, the digital twin system suggested evaluating Melanoma-Associated Antigen A4 (MAGE-A4) and Preferentially Expressed Antigen in Melanoma (PRAME), biomarkers frequently expressed in UCS.^{28,29} Ongoing research explores targeted therapies for these markers. Three relevant clinical trials were accessible within the patient's geographic area. To monitor disease progression, the system recommended continued tracking of serum Cancer Antigen-125 (CA-125) levels based on its established correlation with disease progression identified in the patient's 2021 EHR data.³⁰

Potential treatment trajectories for treatment line four derived from the Digital Twin pipeline are demonstrated in Table 3.

4 Discussion

Extracting meaningful data from unstructured medical text is a prerequisite for precision medicine. In this study, we implemented an LLM-based extraction pipeline to systematically retrieve, structure, and analyze data from real-world EHRs and online sources to support and evaluate diagnostic and targeted therapeutic strategies for constructing patient-specific digital twins for metastatic UCS.

The LLM-based extraction pipeline facilitated timely and accurate synthesis of all relevant full-text scientific publications available through institutional access up to August 15, 2024. The cloud-based LLM achieved accuracy of 0.98 on a complex corpus of medical literature, close to the 0.96 observed by other researchers.³¹ This enabled the generation of evidence-based recommendations and predictive insights grounded in the latest research. Key gaps were observed

in extraction of complex data structures, with recall of 'PFS' (0.77) and 'OS' (0.95) below the overall recall of 0.97. This was due to the fragmented and unstructured way of reporting PFS and OS. Sentences such as "Patient survived for 14 months with the residual tumor post-relapse,"³² make it challenging to accurately determine PFS, as it requires estimation based on prior treatments and the number of treatment cycles. However, estimated PFS may not be accurate if treatment cycles were prolonged. This highlights the challenge of extracting precise outcome data when the primary source lacks comprehensive reporting. We strongly advocate for standards in reporting treatment outcomes, e.g., by clearly stating PFS in months and not date ranges.

In institutional data, unstructured EHR impeded the extraction of key clinical information. This limitation delayed the integration of institutional patient data for informing the management of similar cases. Phase II trials neglecting biomarker-stratification in patients with UCS yielded low objective response rates to ICI therapy, ranging from only zero to ten percent.^{33,34} For our UCS patient, this bottleneck might have precluded ICIs based on high PD-L1 expression, despite the fact that pembrolizumab proved highly efficacious with no adverse effects in this patient. The local LLM system was able to extract structured follow-up data from EHRs across a diverse and complex set of medical documents. While it achieved lower accuracy than the cloud-based model at 0.76, this is in line with the performance of similar models on complex EHR.³⁵ Notably, recall was high across most attributes, with the most critical gap noted in 'biomarkers' at 0.57. The LLM achieved full recall for all biomarkers given as examples for in-context learning, but did not recognize biomarkers that were not explicitly mentioned (e.g., *BRAF* for case four). The LLM again achieved lowest recall for 'PFS' at 0.14. This is due to the highly unstructured and fragmented way of reporting PFS, often across multiple documents.

The European Society for Medical Oncology Precision Medicine Working Group recently established criteria for evaluating the tumor-agnostic potential of molecularly guided therapies, mandating an ORR of $\geq 20\%$ in at least one of five patients across at least four investigated tumor types, with a minimum of five evaluable patients per type.³ Our institutional MTB database identified six analogous cases involving four additional tumor types, most of which exhibited durable responses to ICIs. While the limited number of evaluable patients per tumor type in our single-institution cohort restricted the statistical power, an LLM-driven literature review highlighted an underreporting of PD-L1 expression in UCS in studies conducted to date, despite the known high prevalence of PD-L1 positivity in this malignancy.^{15,16} This underreporting impeded our ability to assess the predictive value of PD-L1 to guide ICI treatment in UCS. Despite the limited sample size of our institutional cohort, the promising outcomes observed suggest that targeting PD-L1 expression in RGT may be a viable therapeutic strategy. The inclusion of diverse tumor types in our institutional cohort further strengthens the role of PD-L1 inhibition in both gynecological and non-gynecological cancers, making it a potential tumor-agnostic marker. Combining PD-1 blockade with bispecific antibodies could offer a promising approach for treating tumors that have not responded to checkpoint inhibitor monotherapy.³⁶

To inform treatment strategies in the event of disease progression, we constructed 21 individualized digital twins, including 7 from our institutional database and 14 from the literature, and queried an LLM-derived database containing 404,265 patient cases. Although our systematic *PubMed* search was specifically limited to the terms "uterine carcinosarcoma" and "endometrial carcinosarcoma," the resulting sample also included other uterine and ovarian malignancies. This is because UCS is frequently reported within the broader context of clinical trials involving more common gynecologic cancers.

The RGT digital twin system generated individualized trajectory predictions for various targeted therapies within a secure local environment that respects patient data privacy, offering guidance on further diagnostics, potential treatment options and continued treatment monitoring with serum CA-125. Additionally, since our real-world cohort comprised only White patients, being able to extract data on patients of other races from a vast corpus of literature helped us validate the generalizability of our treatment recommendations.³⁷

This study successfully demonstrated the utility of RGT digital twins for individualized treatment prediction and response modeling. The digital twin not only provided generalized recommendations for additional diagnostic testing but also incorporated specific clinical details from the patient's treatment history – such as prior pembrolizumab administration – to refine eligibility assessments for targeted therapies. The extraction and analysis of follow-up data revealed that, following the MTB recommendation, the patient received pembrolizumab due to high PD-L1 expression and exhibited a sustained partial response for over 30 months.

This study had several limitations. Firstly, despite a large dataset, the combination of stringent similarity criteria, a limited institutional cohort, and underreported PD-L1 status in published UCS cases prevented us from stratifying patients by PD-L1 status. Efficacy of ICI treatment in PD-L1-positive UCS remains uncertain, and current trials lack PD-L1 as a stratification factor. Bogani et al. listed nine clinical trials currently exploring ICI treatment in UCS, many of which are nearing completion.¹² None of these trials included PD-L1 expression as a stratification factor. Our findings could inform the design of future trials that specifically evaluate ICI efficacy in pMMR UCS with high PD-L1 expression and low to intermediate TMB. Secondly, only somatic biomarkers were included, potentially underestimating

clinical actionability by excluding germline mutations, such as *BRCA1/2*, which are predictive of PARP inhibitor response.³⁸ Thirdly, a local LLM was used for data extraction, impacting extraction performance due to its smaller size and no fine-tuning on German medical texts.³⁹ Lack of a German-language equivalent to the English-language MIMIC labeled medical record dataset⁴⁰ precluded fine-tuning our own model. Finally, the German Network for Personalized Medicine (DNPM) data model is under revision,⁴¹ necessitating the use of a custom data model for this study and highlighting the importance of future validation for compatibility with DNPM v2.

National and international collaborative initiatives, such as the DNPM Data Integration Platform⁴² and the Molecular Tumor Board Portal by Cancer Core Europe,⁴³ aim to enhance MTB decision-making by standardizing and harmonizing data collection across institutions. These platforms stand to benefit substantially from the integration of LLM-based extraction pipelines, which could facilitate the automated extraction of both baseline and follow-up data, thereby enabling the real-time utilization of MTB data across different healthcare systems. Once the DNPM database is fully operational, clinical narratives derived from EHR data could be transformed into HL7 Fast Healthcare Interoperability Resources (FHIR), streamlining interoperability and reducing the biases and costs associated with manual documentation.⁴⁴ Such automation would enable the analysis of larger patient cohorts, thereby providing the statistical power necessary for accurate treatment predictions in rare cancers, a critical step towards advancing personalized oncology. The outcome data could subsequently be used to inform both preclinical research and stratified clinical trials.

Our LLM-enabled precision oncology approach can inform more effective treatment strategies for RGT patients and supports a paradigm shift from organ-based to biology-based tumor classification.

Given the increasing volume and complexity of precision oncology data from MTBs, and the limited availability of precision oncologists to translate this abundance of information into clinically meaningful actions,⁷ there is an urgent need for advanced digital tools to facilitate the extraction, structuring, and analysis of large datasets.¹⁸ Our proof-of-concept study demonstrates the potential of LLMs to efficiently synthesize relevant information for MTB evaluation.

While this study focused on RGTs, the LLM-enabled digital twin approach holds potential for a wide range of refractory cancers. By accurately predicting individual patient trajectories, these digital twins can inform personalized diagnostics and treatment strategies in a timely and cost-effective manner, potentially improving patient outcomes.

Additional information

Ethics statement

Patients from the institutional MTB were included in a clinical registry that allowed for retrospective analyses of clinical and molecular anonymized data in accordance with the Declaration of Helsinki. The retrospective analysis was approved by the Ethics commission of the Medical Faculty of the Technical University of Munich (Reference No. 2023-486-S-SB).

All web-based research procedures were conducted exclusively on publicly accessible, anonymized patient data and in accordance with the Declaration of Helsinki, maintaining all relevant ethical standards.

Acknowledgements

JL is a fellow of the TUM School of Medicine Clinician Scientist Program (project no. H-08). JL acknowledges a 5,000 USD funding in Google Cloud credits from Google's Gemma Academic Research Program. Figures 1 and 2 incorporate icons obtained from Flaticon.com.

Author contributions

Conceptualisation: JL, MT; project administration: JL, MT; resources: JL, NP, KS and SL; validation: JL, NP, LK, KS, SL, US, HB, MT; writing – original draft preparation: JL, MT; writing –review & editing: SM, TD, LM, PM, DF, JNK, DT, LCA, KKB, MB, MK, US, HB. All authors contributed scientific advice and approved the final version of the manuscript.

Funding

JL is supported by the TUM School of Medicine and Health Clinician Scientist Program (project no. H-08). JL receives support through the DKTK School of Oncology Fellowship. PM is funded by the German Federal Ministry of Education and Research through the funding programs Medical Informatics Initiative and National Decade against

Cancer (PM4Onco; 01ZZ2322A). JNK is supported by the German Federal Ministry of Health (DEEP LIVER, ZMV11-2520DAT111; SWAG, 01KD2215B), the Max-Eder-Programme of the German Cancer Aid (grant #70113864), the German Federal Ministry of Education and Research (PEARL, 01KD2104C; CAMINO, 01EO2101; SWAG, 01KD2215A; TRANSFORM LIVER, 031L0312A; TANGERINE, 01KT2302 through ERA-NET Transcan), the German Academic Exchange Service (SECAI, 57616814), the German Federal Joint Committee (Transplant.KI, 01VSF21048) the European Union's Horizon Europe and innovation programme (ODELIA, 101057091; GENIAL, 101096312) and the National Institute for Health and Care Research (NIHR, NIHR213331) Leeds Biomedical Research Centre. DT is funded by the German Federal Ministry of Education and Research (TRANSFORM LIVER, 031L0312A), the European Union's Horizon Europe and innovation programme (ODELIA, 101057091), and the German Federal Ministry of Health (SWAG, 01KD2215B). KKB is supported by the European Union's Horizon Europe and innovation programme (COMFORT, 101079894), Bayern Innovative and Wilhelm-Sander Foundation. MK reports research grants from Sphingotec, Deutsche Krebshilfe, DFG, Senator Roesner Foundation, Dr. Pommer-Jung Foundation, Waltraut Bergmann Foundation, Bavarian State Ministry of Economy, BMBF, G-BA Innovation Fonds. HB reports grants from Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) during the conduct of the study. No other funding is disclosed by any of the authors.

Competing interests

JL received honoraria by the Forum for Continuing Medical Education (FomF) and Novartis for delivering educational lectures. NP has received honoraria for consulting and advisory board participation by AstraZeneca, MSD, GSK, BMS, LabCorp, Illumina, Janssen Cilag, QuIP, and Novartis. DF has received a research grant from OpenAI and holds shares and is an employee at Synagen GmbH. JNK declares consulting services for Owkin, France; DoMore Diagnostics, Norway; Panakeia, UK, and Scailyte, Basel, Switzerland; furthermore JNK holds shares in Kather Consulting, Dresden, Germany; StratifAI GmbH, Dresden, Germany, and Synagen GmbH, Germany, and has received honoraria for lectures and advisory board participation by AstraZeneca, Bayer, Eisai, MSD, BMS, Roche, Pfizer and Fresenius. DT received honoraria for lectures by Bayer and holds shares in StratifAI GmbH, Germany and Synagen GmbH, Germany. KKB has received honoraria for lectures by GE HealthCare and Canon Medical Systems Corporation and serves as advisor for the EU Horizon 2020 project LifeChamps (875329) and the EU IHI Project IMAGIO (101112053). SL has received honoraria for lectures and advisory boards from Taiho Oncology, AstraZeneca, MSD and Janssen-Cilag and provides medical consulting services for NEED Inc. MK has received fees from Springer Press, Biermann Press, Celgene, AstraZeneca, Myriad Genetics, TEVA, Eli Lilly, GSK, Seagen, AllergoSan, FomF, Roche, BESINS, Bayer AG. KS has received honoraria for lectures and advisory boards from BMS, MSD, AstraZeneca, Janssen-Cilag, Roche and Solution akademie GmbH. MK has received honoraria for consulting and advisory board participation by Myriad Genetics, Bavarian KVB, DKMS Life, BLAEK, TEVA, Exeltis, Roche, BESINS, Bayer AG. MK holds shares in AIM GmbH, in-manas GmbH, Therawis Diagnostic GmbH. HB has received personal fees from Roche, AstraZeneca, Gilead, and GlaxoSmithKline outside the submitted work. The authors have no additional financial or non-financial conflicts of interest to disclose.

Data availability statement

Institutional data is available upon reasonable request. All notebooks and prompts used in this study are publicly available at GitHub.

Figures




Figure 1: Workflow from RGT Patient to RGT Digital Twin

RGT Digital Twin is a dynamic system that can integrate diverse data sources to predict individual patient trajectories. Molecular profiling identifies patient biomarkers. LLM capabilities support clinical interpretation of molecular profiles, patient matching to clinical trials, reasoning for cost-coverage requests, medical documentation, and data preparation for advanced computing. Advanced computing techniques such as classification and regression algorithms enable the creation and exploration of Digital Twin models. By adjusting parameters such as biomarker expression or previous treatment strategies, clinicians can model potential patient outcomes and determine suitable treatment strategies. The RGT Digital Twin then integrates outcome data back into the RGT Patient EHR, supporting a continuous learning process.




Figure 2: LLM-enabled RGT Digital Twin Pipeline

To obtain institutional patient data and matching patient profiles from literature, we first filtered institutional records and public data sources (e.g., *PubMed*) by biomarker profiles and primary tumor site. We then extracted structured patient data from EHR using a locally deployed, privacy-preserving LLM, and extracted similar data from published literature using a cloud-based LLM. By utilizing a broader patient population than what is available in institutional data, the RGT Digital Twin system generated personalized treatment plans for MTB evaluation. This method revealed additional treatment options that might have been missed when considering each data source alone.

Tables

Table 1: Evaluation of LLM performance for EHR and literature record extraction

Source File	Data point	Observations	True Positive	True Negative	False Positive	False Negative	Accuracy	Precision	Recall	F1
EHR	Age	7	6	0	1	0	0.86	0.86	1.00	0.86
EHR	Gender	7	7	0	0	0	1.00	1.00	1.00	1.00
EHR	Race	7	7	0	0	0	1.00	1.00	1.00	1.00
EHR	Diagnosis	7	7	0	0	0	1.00	1.00	1.00	1.00
EHR	Biomarkers	7	4	0	0	3	0.57	1.00	0.57	0.73
EHR	Previous treatments	7	2	0	0	5	0.29	1.00	1.00	1.00
EHR	Study treatments	7	7	0	0	0	1.00	1.00	1.00	1.00
EHR	Study treatment response	7	5	0	1	1	0.71	0.83	1.00	0.91
EHR	PFS [months]	7	1	0	0	6	0.14	1.00	0.14	0.25
EHR	OS [months]	7	7	0	0	0	1.00	1.00	1.00	1.00
EHR	TOTAL	70	53	0	2	15	0.76	0.96	0.85	0.91
Source File	Data point	Observations	True Positive	True Negatives	False Positive	False Negatives	Accuracy	Precision	Recall	F1
Literature	Sample size	32	29	3	0	0.00	1.00	1.00	1.00	1.00
Literature	Age	32	29	3	0	0.00	1.00	1.00	1.00	1.00
Literature	Gender	32	7	25	0	0.00	1.00	1.00	1.00	1.00
Literature	Race	32	7	25	0	0.00	1.00	1.00	1.00	1.00
Literature	Diagnosis	32	30	1	0	1.00	0.97	1.00	0.97	0.98
Literature	Biomarkers	32	17	15	0	0.00	1.00	1.00	1.00	1.00
Literature	Previous treatments	32	23	9	0	0.00	1.00	1.00	1.00	1.00
Literature	Study treatments	32	29	2	0	1.00	0.97	1.00	0.97	0.98
Literature	Study treatment response	32	26	5	0	1.00	0.97	1.00	0.96	0.98
Literature	PFS [months]	32	10	19	0	3.00	0.91	1.00	0.77	0.87
Literature	OS [months]	32	18	13	0	1.00	0.97	1.00	0.95	0.97
Literature	TOTAL	352	225	120	0	7.00	0.98	1.00	0.97	0.98

Table 2: ICI treatment outcomes in 7 institutional cases and 14 literature-derived cases

ID	Reference	Diagnosis	Age ^a	Race	PD-L1 Status	TMB (Mut/Mb) ^b	MMR	Addtl. relevant biomarkers (All. frequency)	Treatment Line	ICI treatment (mono/combination)	Response	PFS [months]	OS [months]
1	Institutional	UCS	77	White	CPS: 41, TPS: 3%, IC: 40%	6.3	pMMR (3.6%)	FR α 0.8, PR 0.03, HER2-positive	3	Radiotherapy + pembrolizumab (off-label)	PR	>30 (ongoing)	>132 (ongoing)
2	Institutional	CECSC	37	White	CPS: 75, TPS: 70%, IC: 5%	0	pMMR (1.11%)	None	3	Pembrolizumab (off-label)	PR	>49 (ongoing)	>79 (ongoing)
3	Institutional	CECSC	32	White	CPS: 40, TPS: 40%, IC: <1%	3.1	pMMR (0%)	<i>PIK3CA</i> (p.E545K, 0.26), <i>CHEK2</i> (p.T367Mfs*15, 0.79)	2	Pembrolizumab (off-label)	PD	1	15 (deceased)
4	Institutional	CECSC	85	White	CPS: 81, TPS: 80%, IC: 1%	11	pMMR (4.6%)	<i>BRAF</i> (p.D594N, 0.27), <i>KMT2C</i> (p.Q192Tfs*28, 0.29)	4	Pembrolizumab (off-label)	PR, PD	18	72 (deceased)
5	Institutional	CEAD	37	White	CPS: 95, TPS: 90%, IC: 5%	5.5	pMMR (4.6%)	None	2	Ipilimumab/nivolumab, nivolumab maintenance (off-label)	CR	>45	>45 (ongoing)
6	Institutional	USC	61	White	CPS: 40, TPS: 30%, IC: 8%	13.4	pMMR (1.89%)	<i>PIK3CA</i> (p.E545K, 0.06), <i>PTEN</i> (p.K128Rfs*6, 0.13), <i>PTEN</i> (p.Y240delins*, 0.06), FR α : 0%, HER2: Score 0, Trop2: 100%	3	Pembrolizumab + lenvatinib (in-label)	PD	3	>69 (ongoing)
7	Institutional	Undifferentiated Sarcomatoid Carcinoma of the Pancreas	60	White	CPS: 85, TPS: 80%, IC: 4%	3.2	pMMR (2.61%)	<i>KRAS</i> (p.G12C; 0.38)	3	Pembrolizumab (off-label)	PR, PD	6	19 (deceased)
8	PMID: 32620662	UCS	65	Asian (Japanese)	positive	n/a	dMMR/MSI-H	None	2	Radiotherapy + pembrolizumab	CR, PD	10	16 (deceased)
9	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		3	Pembrolizumab + lenvatinib	PD	3.3	9.9 (deceased)
10	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		3	Pembrolizumab + lenvatinib	PD	0.9	2.8 (deceased)
11	PMID: 34401435	UCS	n/a ^c	n/a ^c	positive	n/a	dMMR/MSI-H		3	Pembrolizumab + lenvatinib	PD	1.6	2.4 (deceased)
12	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		3	Pembrolizumab + lenvatinib	PD	2.6	2.8 (deceased)
13	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		5	Pembrolizumab + lenvatinib	PD	1.9	2.1 (deceased)
14	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		4	Pembrolizumab + lenvatinib	SD	- (ongoing)	4.4 (alive at data cut-off)
15	PMID: 34401435	UCS	n/a ^c	n/a ^c	negative	n/a	pMMR		3	Pembrolizumab + lenvatinib	SD, PD	11.2	12.6 (alive at data cut-off)
16	PMID: 29386312	UCS	55	n/a	1+, low positive	169	pMMR	<i>POLE</i> -mutated	4	Pembrolizumab	PR	>12 (ongoing)	39 (alive at data cut-off)
17	PMID: 38881561	UCS	68	n/a	n/a	6	pMMR	<i>PTEN</i> K128T, <i>ESR1</i> -amplified (8/8 exons, est. 11 copies), ER positive	2	Pembrolizumab + lenvatinib + letrozole	PR	>36 (ongoing)	45 (alive at data cut-off)
18	PMID: 30442730	UCS	59	n/a	n/a	n/a	n/a		2	Pembrolizumab	MR	4	n/a
19	PMID: 33004543	UCS	66	Asian (Japanese)	n/a	n/a	dMMR	Highly predisposing HLA haplotype for narcolepsy	3	Pembrolizumab	PD	2	Deceased, 72 days post pembrolizumab, OS n/a
20	PMID: 31149529	UCS	68	n/a	n/a	n/a	n/a		2	PD-1 antibody + CTLA-4 antibody	PR	>5 (ongoing)	N/a, alive
21	PMID: 35434237	UCS	62	n/a	n/a	14	pMMR	Germline <i>NBN</i> mutation, (c.2117C>G, p.Ser706Ter) HER2-low (Score 1+)	3	Avelumab + axetinib	PR	>15 (ongoing)	48 (alive at data cut-off)

a. Current age at data cut-off (publication)

b. TMB: <5: low, 5-15: intermediate, \geq 15: high

c. Patients 9-15: Median age: 63 (range: 58-64), White: 3, Black: 4. Individual data for age & race not reported.

PMID: PubMed-ID

Note: 'n/a' entries indicate data not available for the specific case.

Table 3: Digital twin pipeline provided the following individualized treatment targets for case 1

Biomarker	Prevalence in UCS	Biomarker Result	Treatment	Expected treatment response	Reference
HER2	Expressed in 1/3 of UCS ^{27,45}	Positive (Histopathological report, January, 2021 "HER2 is positive in some of the tumor cells with an intermediate level of positivity and an incomplete level of circumferential positivity.")	Trastuzumab deruxtecan T-DM1	The phase II STATICE trial enrolled 22 HER2-high and 10 HER2-low patients with recurrent UCS. Objective response rates (ORRs) were 54.5% and 70% in the HER2-high and HER2-low groups, respectively. Median progression-free survival (PFS) was 6.2 months for HER2-high patients and 13.3 months for HER2-low patients. Overall survival (OS) was 6.7 months for HER2-high patients and not reached for HER2-low patients. Three patients in each group had received at least three prior lines of therapy. T-DM1 demonstrated significant antitumor activity in HER2-overexpressing CS xenograft models, resulting in prolonged survival compared to trastuzumab.	Phase II study ²⁶ Preclinical evidence ⁴⁶
ER	Estrogen receptor: 20–30%; progesterin receptor: 5–40% ¹²	80%	Anihormonal treatment (e.g., anastrozole)	The phase II PARAGON study enrolled seven patients with UCS and evaluated anastrozole treatment. A clinical benefit rate (CBR) of 43% was observed at three months, with a median duration of clinical benefit of 5.6 months. While stable disease was noted in three patients, no objective responses were achieved. Median progression-free survival was 2.7 months. 43% of the entire cohort (UCS and leiomyosarcoma) had received prior chemotherapy.	Phase II study ⁴⁷
	<i>ESR1</i> Amplification	7%	Pembrolizumab + lenvatinib + letrozole	A patient with metastatic, pMMR, and <i>ESR1</i> -amplified UCS achieved a durable partial response of 36 months with third-line treatment combining pembrolizumab, lenvatinib, and letrozole.	Case report ⁴⁸
FR-alpha	Expressed in 1/3 of UCS ⁴⁹	Not determined.	Mirvetuximab soravtansine + pembrolizumab	Patients with FR-alpha positive tumors may be eligible for combination therapy with mirvetuximab soravtansine and pembrolizumab. This approach is supported by results from a phase II trial (NCT03835819) demonstrating efficacy in pMMR-positive patients, including those with prior pembrolizumab treatment failure. Interim analysis in endometrial cancers showed ORR of 37.5%. No stratified analysis for UCS available.	Phase II study ⁵⁰
HRD	Unknown	Not determined.	Poly (ADP-Ribose) Polymerase Inhibitor (PARPi)	UCS cell lines exhibiting HRD signature demonstrated significantly increased sensitivity to olaparib compared to homologous recombination proficient UCS cell lines, both in vitro and in vivo.	Preclinical evidence ⁵¹
MAGE-A4	Expressed in 91% of carcinosarcomas ²⁸	Not determined.	Bispecific T Cell Engaging Receptor Molecule targeting MAGE-A4/8 expression	Patients with MAGE-A4-positive UCS may be eligible for participation in an ongoing clinical trial located in Bavaria: A Phase Ia/Ib First-In-Human Clinical Trial to Evaluate the Safety, Tolerability and Initial Anti-tumor Activity of IMA401, a Bispecific T Cell Engaging Receptor Molecule (TCER®), in Patients With Recurrent and/or Refractory Solid Tumors.	https://clinicaltrials.gov/study/NCT05359445
PRAME	Expressed in 60% of UCS ^{29,52}	Not determined.	Bispecific T Cell-Engaging Receptor Molecule targeting PRAME Genetically Modified Autologous T Cells Expressing a T Cell Engaging Receptor Recognizing PRAME as Monotherapy or in Combination with Nivolumab	Patients with PRAME-positive UCS may be eligible for participation in two ongoing clinical trials located in Bavaria: a) IMA402-101: A Phase I/II First-In-Human Clinical Trial to Evaluate the Safety, Tolerability and Anti-Tumor Activity of IMA402, a Bispecific T Cell-Engaging Receptor Molecule (TCER) Targeting PRAME, in Patients With Recurrent and/or Refractory Solid Tumors. b) IMA203-101: Phase 1 Study Evaluating Genetically Modified Autologous T Cells Expressing a TCR Recognizing a Cancer/Germine Antigen as Monotherapy or in Combination With Nivolumab in Patients With Recurrent and/or Refractory Solid Tumors	a) https://clinicaltrials.gov/study/NCT05958121 b) https://clinicaltrials.gov/study/NCT03686124
Trop2	Expressed in 1/3 of UCS ^{53,54}	Not determined.	Sacituzumab govitecan	Twenty-two patients with Trop2-positive recurrent endometrial cancer were enrolled in a phase II study evaluating sacituzumab govitecan. Of these, three patients had UCS. Among the 20 response-evaluable patients, an objective response rate of 35% was observed. Median progression-free survival (PFS) and overall survival (OS) were 5.7 months and 22.5 months, respectively. The median (range) number of prior anticancer regimens was 3 (1–6). Sacituzumab govitecan demonstrated significant tumor growth inhibition and improved 90-day overall survival in Trop2-positive carcinosarcoma cell lines compared to Trop2-negative controls.	Phase II study ⁵⁵ Preclinical evidence ⁵⁴
Serum CA-125	Elevated serum CA-125 levels in 62% of UCS patients with FIGO stage III or IV ³⁰	Elevated at progression in 2021, normalized since 2022 correlating with partial response.	Treatment monitoring with Serum CA-125	Preoperative CA-125 elevation correlates with extrauterine disease and deep myometrial invasion in patients with UCS. Postoperatively, elevated CA-125 is an independent prognostic indicator of poor survival. These findings suggest that CA-125 could serve as a valuable serum marker for managing UCS patients.	Retrospective analysis ³⁰


References

- [1] G. Gatta et al. “Rare cancers are not so rare: the rare cancer burden in Europe”. In: *Eur. J. Cancer* 47.17 (Nov. 2011), pp. 2493–2511.
- [2] A. Lainé, B. Hanvic, and I. Ray-Coquard. “Importance of guidelines and networking for the management of rare gynecological cancers”. In: *Curr. Opin. Oncol.* 33.5 (Sept. 2021), pp. 442–446.
- [3] C. Westphalen et al. “The ESMO Tumour-Agnostic Classifier and Screener (ETAC-S): a tool for assessing tumour-agnostic potential of molecularly guided therapies and for steering drug development”. In: *Annals of Oncology* (Aug. 2024).
- [4] P. Horak et al. “Comprehensive Genomic and Transcriptomic Analysis for Guiding Therapeutic Decisions in Patients with Rare Cancers”. In: *Cancer Discov.* 11.11 (Nov. 2021), pp. 2780–2795.
- [5] J. A. Ledermann. “PARP inhibitors in ovarian cancer”. In: *Ann. Oncol.* 27 Suppl 1 (Apr. 2016), pp. i40–i44.
- [6] F. André et al. *Forget lung, breast or prostate cancer: why tumour naming needs to change*. en. <http://dx.doi.org/10.1038/d41586-024-00216-3>. Accessed: 2024-8-28. Jan. 2024.
- [7] A. M. Tsimberidou et al. “Molecular tumour boards - current and future considerations for precision oncology”. In: *Nat. Rev. Clin. Oncol.* 20.12 (Dec. 2023), pp. 843–863.
- [8] T. Botsis et al. “Precision Oncology Core Data Model to Support Clinical Genomics Decision Making”. In: *JCO Clin Cancer Inform* 7 (Apr. 2023), e2200108.
- [9] J. N. Acosta et al. “Multimodal biomedical AI”. In: *Nat. Med.* 28.9 (Sept. 2022), pp. 1773–1784.
- [10] M. N. Kamel Boulos and P. Zhang. “Digital Twins: From Personalised Medicine to Precision Public Health”. In: *J Pers Med* 11.8 (July 2021).
- [11] U.S. Food and Drug Administration. “Paving the way for personalized medicine: FDA’s role in a new era of medical product development”. In: *Silver Spring, MD: US Food and Drug Administration* (2013).
- [12] G. Bogani et al. “Endometrial carcinosarcoma”. In: *Int. J. Gynecol. Cancer* 33.2 (Feb. 2023), pp. 147–174.
- [13] E. Katsoulakis et al. “Digital twins for health: a scoping review”. In: *NPJ Digit Med* 7.1 (Mar. 2024), p. 77.
- [14] L. C. Adams et al. “Leveraging GPT-4 for Post Hoc Transformation of Free-text Radiology Reports into Structured Reporting: A Multilingual Feasibility Study”. In: *Radiology* 307.4 (May 2023), e230725.
- [15] S. Hacking et al. “Landscape of Immune Checkpoint Inhibition in Carcinosarcoma (MMMT): Analysis of IDO-1, PD-L1 and PD-1”. In: *Pathol. Res. Pract.* 216.4 (Apr. 2020), p. 152847.
- [16] T. M. Jenkins et al. “PD-L1 and Mismatch Repair Status in Uterine Carcinosarcomas”. In: *Int. J. Gynecol. Pathol.* 40.6 (Nov. 2021), pp. 563–574.
- [17] U. Kucukgoz Gulec et al. “Prognostic significance of programmed death-1 (PD-1) and programmed death-ligand 1 (PD-L1) expression in uterine carcinosarcoma”. In: *Eur. J. Obstet. Gynecol. Reprod. Biol.* 244 (Jan. 2020), pp. 51–55.
- [18] The Lancet Oncology. “Incorporating whole-genome sequencing into cancer care”. In: *Lancet Oncol.* 25.8 (Aug. 2024), p. 945.
- [19] T. Hegghammer. “OCR with Tesseract, Amazon Textract, and Google Document AI: a benchmarking experiment”. In: *Journal of Computational Social Science* 5.1 (May 2022), pp. 861–882.
- [20] Gemma Team et al. “Gemma 2: Improving Open Language Models at a Practical Size”. In: (July 2024). arXiv: 2408.00118 [cs.CL].
- [21] S. M. Xie et al. “An Explanation of In-context Learning as Implicit Bayesian Inference”. In: (Nov. 2021). arXiv: 2111.02080 [cs.CL].
- [22] Gemini Team et al. “Gemini 1.5: Unlocking multimodal understanding across millions of tokens of context”. In: (Mar. 2024). arXiv: 2403.05530 [cs.CL].
- [23] E. Mosqueira-Rey et al. “Human-in-the-loop machine learning: a state of the art”. In: *Artificial Intelligence Review* 56.4 (Apr. 2023), pp. 3005–3054.
- [24] I. J. Marshall and B. C. Wallace. “Toward systematic review automation: a practical guide to using machine learning tools in research synthesis”. In: *Syst. Rev.* 8.1 (July 2019), p. 163.
- [25] ClinicalTrials.gov. *A Phase 2 Study of Mirvetuximab Soravtansine (IMGN853) and Pembrolizumab in Endometrial Cancer (EC) (NCT03835819)*. en. <https://clinicaltrials.gov/study/NCT03835819>. Accessed: 2024-8-16. 2024.
- [26] T. Nishikawa et al. “Trastuzumab Deruxtecan for Human Epidermal Growth Factor Receptor 2-Expressing Advanced or Recurrent Uterine Carcinosarcoma (NCCH1615): The STATICE Trial”. In: *J. Clin. Oncol.* 41.15 (May 2023), pp. 2789–2799.

- [27] H. Yoshida et al. “Discordances in expression of human epidermal growth factor receptor 2 between primary and metastatic uterine carcinosarcoma: A proposal for HER2-targeted therapy specimen selection”. In: *Ann. Diagn. Pathol.* 65 (Aug. 2023), p. 152150.
- [28] M. B. Resnick et al. “Cancer-testis antigen expression in uterine malignancies with an emphasis on carcinosarcomas and papillary serous carcinomas”. In: *Int. J. Cancer* 101.2 (Sept. 2002), pp. 190–195.
- [29] A. Alrohaibani et al. “Preferentially Expressed Antigen in MELanoma Expression in Uterine and Ovarian Carcinosarcomas”. In: *Int. J. Gynecol. Pathol.* 43.3 (May 2024), pp. 284–289.
- [30] G. S. Huang et al. “Serum CA125 predicts extrauterine disease and survival in uterine carcinosarcoma”. In: *Gynecol. Oncol.* 107.3 (Dec. 2007), pp. 513–517.
- [31] A. Konet et al. “Performance of two large language models for data extraction in evidence synthesis”. In: *Res Synth Methods* (June 2024).
- [32] M. Yano et al. “Pembrolizumab and Radiotherapy for Platinum-refractory Recurrent Uterine Carcinosarcoma With an Abscopal Effect: A Case Report”. In: *Anticancer Res.* 40.7 (July 2020), pp. 4131–4135.
- [33] M. M. Rubinstein et al. “Durvalumab with or without tremelimumab in patients with persistent or recurrent endometrial cancer or endometrial carcinosarcoma: A randomized open-label phase 2 study”. In: *Gynecol. Oncol.* 169 (Feb. 2023), pp. 64–69.
- [34] S. Lheureux et al. “Translational randomized phase II trial of cabozantinib in combination with nivolumab in advanced, recurrent, or metastatic endometrial cancer”. In: *J Immunother Cancer* 10.3 (Mar. 2022).
- [35] K. Saab et al. “Capabilities of Gemini Models in Medicine”. In: (Apr. 2024). arXiv: 2404.18416 [cs.AI].
- [36] M. Merz. “A comeback for checkpoint inhibition in multiple myeloma”. In: *Nature Cancer* (2024), pp. 1–3.
- [37] K. Arora et al. “Genetic Ancestry Correlates with Somatic Differences in a Real-World Clinical Cancer Sequencing Cohort”. In: *Cancer Discov.* 12.11 (Nov. 2022), pp. 2552–2565.
- [38] L. B. Mendes Gomes et al. “Primary Peritoneal Carcinosarcoma in a Breast Cancer Patient Harboring a Germline BRCA2 Pathogenic Variant: Case Report”. In: *Case Rep. Oncol.* 17.1 (Jan. 2024), pp. 1–9.
- [39] T. Han et al. “MedAlpaca – An Open-Source Collection of Medical Conversational AI Models and Training Data”. In: (Apr. 2023). arXiv: 2304.08247 [cs.CL].
- [40] A. E. W. Johnson et al. “MIMIC-IV, a freely accessible electronic health record dataset”. In: *Sci Data* 10.1 (Jan. 2023), p. 1.
- [41] B. Siebrasse. *MTB Kerndatensätze_V01 (öffentlich zugänglich) - DNPM AG - MedInf - IBMI-UT Confluence*. <https://ibmi-ut.atlassian.net/wiki/spaces/DAM/pages/2592722/>. Accessed: 2024-8-16. 2024.
- [42] A. L. Illert et al. “The German Network for Personalized Medicine to enhance patient care and translational research”. In: *Nat. Med.* 29.6 (June 2023), pp. 1298–1301.
- [43] D. Tamborero et al. “The Molecular Tumor Board Portal supports clinical decisions and automated reporting for precision oncology”. In: *Nat Cancer* 3.2 (Feb. 2022), pp. 251–261.
- [44] Li Yikuan et al. “FHIR-GPT Enhances Health Interoperability with Large Language Models”. In: *NEJM AI* 1.8 (July 2024), A1cs2300301.
- [45] T. M. Jenkins et al. “HER2 Overexpression and Amplification in Uterine Carcinosarcomas With Serous Morphology”. In: *Am. J. Surg. Pathol.* 46.4 (Apr. 2022), pp. 435–442.
- [46] R. Nicoletti et al. “T-DM1, a novel antibody-drug conjugate, is highly effective against uterine and ovarian carcinosarcomas overexpressing HER2”. In: *Clin. Exp. Metastasis* 32.1 (Jan. 2015), pp. 29–38.
- [47] R. J. Edmondson et al. “Phase 2 study of anastrozole in rare cohorts of patients with estrogen receptor/progesterone receptor positive leiomyosarcomas and carcinosarcomas of the uterine corpus: The PARAGON trial (ANZGOG 0903)”. In: *Gynecol. Oncol.* 163.3 (Dec. 2021), pp. 524–530.
- [48] J. L. Soiffer et al. “Durable partial response to pembrolizumab, lenvatinib, and letrozole in a case of recurrent uterine carcinosarcoma with ESR1 gene amplification”. In: *Gynecol Oncol Rep* 54 (Aug. 2024), p. 101426.
- [49] A. Saito et al. “Folate receptor alpha is widely expressed and a potential therapeutic target in uterine and ovarian carcinosarcoma”. In: *Gynecol. Oncol.* 176 (Sept. 2023), pp. 115–121.
- [50] R. L. Porter et al. “Abstract CT008: A phase 2, two-stage study of mirvetuximab soravtansine (IMGN853) in combination with pembrolizumab in patients with microsatellite stable (MSS) recurrent or persistent endometrial cancer”. In: *Cancer Res.* 84.7_Supplement (Apr. 2024), CT008–CT008.
- [51] J. R. Tymon-Rosario et al. “Homologous recombination deficiency (HRD) signature-3 in ovarian and uterine carcinosarcomas correlates with preclinical sensitivity to Olaparib, a poly (adenosine diphosphate [ADP]-ribose) polymerase (PARP) inhibitor”. In: *Gynecol. Oncol.* 166.1 (July 2022), pp. 117–125.
- [52] J. Roszik et al. “Overexpressed PRAME is a potential immunotherapy target in sarcoma subtypes”. In: *Clin. Sarcoma Res.* 7 (June 2017), p. 11.

- [53] R. Raji et al. “Uterine and ovarian carcinosarcomas overexpressing Trop-2 are sensitive to hRS7, a humanized anti-Trop-2 antibody”. In: *J. Exp. Clin. Cancer Res.* 30.1 (Nov. 2011), p. 106.
- [54] S. Lopez et al. “Preclinical activity of sacituzumab govitecan (IMMU-132) in uterine and ovarian carcinosarcomas”. In: *Oncotarget* 11.5 (Feb. 2020), pp. 560–570.
- [55] A. Santin et al. “Preliminary results of a phase II trial with sacituzumab govitecan-hziy in patients with recurrent endometrial carcinoma overexpressing Trop-2”. In: *J. Clin. Orthod.* 41.16_suppl (June 2023), pp. 5599–5599.
- [56] Mirza Mansoor R. et al. “Dostarlimab for Primary Advanced or Recurrent Endometrial Cancer”. In: *N. Engl. J. Med.* 388.23 (June 2023), pp. 2145–2158.
- [57] S. N. Westin et al. “Durvalumab Plus Carboplatin/Paclitaxel Followed by Maintenance Durvalumab With or Without Olaparib as First-Line Treatment for Advanced Endometrial Cancer: The Phase III DUO-E Trial”. In: *J. Clin. Oncol.* 42.3 (Jan. 2024), pp. 283–299.
- [58] N. Colombo et al. “Atezolizumab and chemotherapy for advanced or recurrent endometrial cancer (AtTEnd): a randomised, double-blind, placebo-controlled, phase 3 trial”. In: *Lancet Oncol.* (Aug. 2024).
- [59] N. S. Azad et al. “Nivolumab Is Effective in Mismatch Repair-Deficient Noncolorectal Cancers: Results From Arm Z1D-A Subprotocol of the NCI-MATCH (EAY131) Study”. In: *J. Clin. Oncol.* 38.3 (Jan. 2020), pp. 214–222.
- [60] J. A. How et al. “Toxicity and efficacy of the combination of pembrolizumab with recommended or reduced starting doses of lenvatinib for treatment of recurrent endometrial cancer”. In: *Gynecol. Oncol.* 162.1 (July 2021), pp. 24–31.

Supplements



Supplementary Figure 1: The previous treatment regime did not mention the addition of dostarlimab as the consensus statement was published before EMA approval of dostarlimab on October 12, 2023. Reported response rates and median PFS with third-line therapy were 5.5% and 1.8 months, respectively. The 5-year overall survival rate has not changed in decades (31.9% in 1975 to 33.8% in 2012)¹². The targeted treatment approach could result in a prolongation of PFS and consequently a better OS.

Supplementary Table 1: Detailed results of MTB for case 1

Submitting hospital	Block material	Biomarkers tested before MTB	Tumor cell count	Panel used	Single Nucleotide Variants (SNV)/Indel	Copy number changes	Splice variations/translocations
TUM University Hospital, Munich, Bavaria, Germany	Supracervical lymph node metastasis (2021)	ER: 80% PR: 3% HER2: Positive	60%	TSO 500 (DNA)/TST 170 (RNA)	None detected	None detected	None detected
Gene	Reference number	Exon	cDNA	Protein	Allele frequency	COSMIC database v90	Class
<i>TP53</i>	NM_000546	4	c.150delT	p.150Mfs*73	14%	COSV52758078	4
<i>PPP2R1A</i>	NM_014225	6	c.771G>T	p.W257C	12%	COSV59043009	4
<i>NOTCH4</i>	NM_004557	18	c.2780G>A	p.C927Y	8%	No entry	3
<i>RUNX1</i>	NM_001754	9	c.1070C>T	p.P357L	9%	No entry	3
<i>AR</i>	NM_000044	1	c.476C>G	p.A159G	7%	No entry	3

Supplementary Table 2: Data structure for RGT Digital Twin

Attribute	Data Type	Description
n	Integer	Number of patients in the study
age	String	Age of patients in the study; string as many studies contain ranges
gender	String	Gender(s) of patients in the study
race	String	Race(s) of patients in the study
diagnosis	String	Diagnosis of patients in the study
biomarkers	{ 'pd-l1': String, 'tmb/mb': String, 'msi/mss':String, 'others': String }	Biomarkers (e.g., PD-L1) determined and discussed in the study; returned as dictionary for simplified analysis
previous treatments	String	Description of previous treatments (and response)
study treatment	String	Treatments discussed in the study
study treatment response	{ 'treatment response': String, 'adverse effects': String }	Response to treatments discussed in the study including adverse effects; returned as dictionary for simplified analysis
PFS	String	Progression-free survival (PFS) in months reported for study treatment; string as many studies contain ranges
OS	String	Overall survival (OS) in months reported for patient cases; string as many studies contain ranges

Supplementary Table 3: Baseline characteristics of institutional MTB cases at TUM University Hospital

ID	Gender	Primary site	Age at diagnosis	Stage at diagnosis	Sites of metastases	Time of MTB	Site sequenced	Tumor cell count	PD-L1	TMB (Mut/Mb)	MMR
1	female	Uterine Carcinosarcoma (UCS)	66	FIGO IIIC2, pT3a, pN2 (3/67), R0	Cervical lymph node metastasis, pelvic recurrence, retroperitoneal lymph node metastases	2021	Cervical lymph node metastasis	60%	CPS: 41 TPS: 3% IC: 40%	6.3 (intermediate)	pMMR/MSS (3.6%)
2	female	Cervical Squamous Cell Carcinoma (CESC)	30	FIGO IB1, pT1b1, pNX, G3, R0	Pelvic recurrence, liver metastasis	2020	Primary surgery	70%	CPS: 75 TPS: 70% IC: 5%	0 (low)	pMMR/MSS (1.11%)
3	female	Cervical Squamous Cell Carcinoma (CESC)	31	FIGO IVB, cT2a, pN1 (12/94), cM1 (PER)	Peritoneal metastases	2020	Peritoneal metastasis	75%	CPS: 40 TPS: 40% IC: <1%	3.1 (low)	pMMR/MSS (0%)
4	female	Cervical Squamous Cell Carcinoma (CESC)	79	FIGO IVa, cT2b2, cN1, cM0, G2	Lymph node metastases	2021	Primary tumor biopsy	80%	CPS: 81 TPS: 80% IC: 1%	11 (intermediate)	pMMR/MSS (4.6%)
5	female	Cervical Adenocarcinoma (CEAD)	32	FIGO IVB, cT2b, cN1, cM1 (LYM)	Lymph node metastases	2021	Ileocecal resection	40%	CPS: 95 TPS: 90% IC: 5%	5.5 (intermediate)	pMMR/MSS (3.28%)
6	female	Uterine Serous Carcinoma (USC)	55	FIGO IA1, pT1a, pNx, L0, V0, Pn0, R0	Peritoneal metastases, lymph node metastases	2024	Inguinal lymph node metastasis	30%	CPS: 40 TPS: 30% IC: 8%	13.4 (intermediate)	pMMR/MSS (1.89%)
7	male	Undifferentiated Sarcomatoid Carcinoma of the Pancreas	59	pT3, pN1 (3/81), L1, V1, Pn1, R0	Lcooregional recurrence, liver metastases	2022	Liver metastasis	70%	CPS: 85 TPS: 80% IC: 4%	3.2 (low)	pMMR/MSS (2.61%)

Supplementary Table 4: Seven studies on ICI treatment in UCS lacked patient-level data necessary for individual digital twin creation

Trial	Recruitment period	Phase	Experimental group	Control group	Sample size	Treatment response	Median follow-up (months)	Median PFS (months) experimental vs control
RUBY trial ⁵⁶	2019-2021	III	Carboplatin/paclitaxel (CP) + dostarlimab x 6 cycles + maintenance with dostarlimab	CP + placebo x 6 cycles + maintenance with placebo	Overall: 494UCS: n = 44 dMMR: 118(UCS: 5) pMMR: 376(UCS: 39)	Not stratified for UCS Overall: Hazard Ratio (HR) = 0.64 (0.51-0.80) p <0.001 dMMR: HR = 0.28 (0.16-0.50) p <0.001 pMMR: HR = 0.76 (0.59-0.98) Overall: HR = 0.71 (0.57-0.89) p = 0.003	Not stratified for UCS Overall: 25.4 dMMR: 24.8 pMMR NA	NA
DUO-E trial ⁵⁷	2020-2022	III	CP + durvalumab x 6 cycles + maintenance with durvalumab	CP + placebo x 6 cycles + maintenance with placebo	Overall 479 UCS: n = 61 dMMR 95 pMMR 384	p = 0.003 dMMR: HR = 0.42 (0.22-0.80) pMMR: HR = 0.77 (0.60-0.97) Histology: other, including UCS (27/39) HR = 0.76 (0.46-1.25), n.s. Overall: HR = 0.74 (0.61-0.91), p = 0.02 UCS: HR = 0.88 (0.45-1.73), n.s.	Not stratified for UCS Control 12.6 Experimental 15.4 dMMR 10.2 pMMR 12.8	Not stratified for UCS Overall 10.2 vs 9.6 dMMR NR vs 7 pMMR 9.9 vs. 9.7
AtTEnd trial ⁵⁸	2018-2022	III	CP + atezolizumab x 6 cycles + maintenance with atezolizumab	CP + placebo x 6 cycles + maintenance with placebo	Overall: 549 UCS: n = 50 dMMR: 125 pMMR: 409	UCS: HR = 0.88 (0.45-1.73), n.s. dMMR: HR = 0.36 (0.23-0.57), p = 0.0005 UCS: HR = 0.41 (0.03-6.62), n.s. pMMR: HR = 0.92 (0.73-1.16), n.s. UCS: not specified	Not stratified for UCS Overall: 28.3 dMMR: 26.2 pMMR NA	Not stratified for UCS Overall: 10.1 vs 8.9 dMMR NR vs. 6.9 pMMR 9.5 vs 9.2
Single-center, randomized, open-label, phase II trial ³³	N/a Data cut-off: December 2021	II	Arm 1: Durvalumab Arm 2: Durvalumab + tremelimumab	None	Overall: 82 UCS: 16 Arm 1: 6 Arm 2: 10	Overall: Arm 1 Overall Response Rate (ORR): 10.8% Arm 2 ORR: 5.3% UCS: ORR: 0%	N/a	N/a
NCI-MATCH (EAY131) ⁵⁹	2016-2017	II	Nivolumab	None	Overall: 42 UCS: n = 4	Not stratified for UCS Overall ORR: 36%	17.3	Not stratified for UCS 6-month PFS rate: 51.3% 12-month PFS rate: 46.2% 18-month PFS rate: 31.4% Not stratified for UCS
Retrospective institutional analysis from The University of Texas MD Anderson Cancer Center ⁶⁰	2019-2020	Retrospective study of institutional data	Pembroliuzumab + lenvatinib Recommended dose of lenvatinib vs. reduced dose of lenvatinib	None	Overall: n = 61 Recommended dose: n = 14 Reduced dose: n = 47 UCS: n = 16 Recommended dose: n = 3 Reduced dose: n = 13	ORR: Overall: 36.1% UCS: 25% (3/12) Clinical benefit rate (CBR): Overall: 68.9% UCS: 58.3% (7/12)	Not stratified for UCS Overall: Recommended dose: 3.2 Reduced dose: 5.5	Not stratified for UCS Recommended dose: 8.6 Reduced dose: 9.4
Multicenter, randomized, phase II trial ⁵⁴	2018-2019	II	Cabozantinib + nivolumab	None	Arm A: 36 Arm B: 18 Arm C: UCS: n = 10 pMMR: 100%	Arm C (UCS): ORR: 10% 1 PR, 5 SD	Overall (Arm A, B, C): 15.9	Arm C (UCS): Median SD duration: 3.2 (range 2.8-7.6)