

DIAGONAL REPRESENTATION OF OPEN STRING STAR AND MOYAL PRODUCT

D.M. Belov*

Department of Physics
Rutgers University
136 Frelinghuysen Rd.
Piscataway, NJ 08854, USA
belov@physics.rutgers.edu

Abstract

We explicitly find the spectrum of the operators M^{rs} and \widetilde{M}^{rs} , which specify the \star -product in the matter and ghost sectors correspondingly. Further we derive the diagonal representation for the 3-string vertices in both sectors. Using this representation we identify the appearing Moyal structures in the matter sector. In addition to the continuous non-commutativity parameter $\theta(\kappa)$ found in [6] we find the discrete non-commutativity parametrized by θ_ξ .

*On leave from Steklov Mathematical Institute, Moscow, Russia.

Contents

1	Introduction	3
2	Review of the 3-string vertex	3
3	Spectrum of the operators CU and UC (matter)	5
3.1	Notations	5
3.2	Eigenvectors of the operators CU and UC	7
3.3	Summary	9
4	Spectrum of the operators defining ghost vertex	11
4.1	Spectrum of the operators $C\tilde{U}$ and $\tilde{U}C$ (ghosts)	11
4.2	Spectrum of the operators $C'\tilde{U}'$ and $\tilde{U}'C'$ (ghosts)	12
5	Diagonal representation of the open string star	13
5.1	Matter 3-vertex	13
5.2	Ghost 3-vertex	14
6	Moyal structure	16
6.1	Oscillator vertex for the Moyal product	16
6.2	Identification of Moyal structures	18
7	Discussion	18
	Appendix	21
A	Normalization of the coordinate and momentum eigenstates	21
B	Conformal Ghosts	22
B.1	Search for the discrete spectrum	22
B.1.1	Parametrization of the eigenvalues	22
B.1.2	Computation of the integral	22
B.1.3	Solving of the equation	24
B.2	Generating function for discrete spectrum	25
B.3	Normalization of the vectors from the discrete spectrum	26
B.4	Generating function for the continuous spectrum	27
B.4.1	Calculation of $A^{(\kappa)}$	27
B.4.2	Generating function	28
B.5	Normalization of the vectors from continuous spectrum	29

1 Introduction

Recently it has been a resurgence of interest in string field theory (SFT) [1]. Along with the applications to such problems as the tachyon condensation [2], [3, 4, 5] the intrinsic structure of SFT is also investigated. In particular recently M. Douglas, H. Liu, G. Moore and B. Zwiebach [6] have shown that open zero-momentum string field algebra can be represented as a continuous product of mutually commuting Moyal algebras. The first attempt to describe the open string star product as a series of Moyal products was made in [7]. The aim of the present paper is to generalize these results to the matter sector including zero modes and to the ghost sector. One hopes that finding such a diagonal representation will lead to a better understanding of string field \star -algebra structure and finding the solutions of SFT equations of motion.

The paper is organized as follows. In Section 2 we review the structure of 3-string vertex in the matter and ghost sectors. In Section 3 we find the spectrum of operators CU and UC defining the Neumann matrices in the matter sector. Subsection 3.3 contains summary on the spectrum of the matter operator CU . In Section 4 we consider the spectrum of various operators defining the ghost Neumann matrices. In Section 5 we derive the diagonal representation for the matter and ghost 3-string vertices. In Section 6 identify the Moyal structures appearing in the diagonal representation of the vertices.

Appendices contain necessary technical information. In Appendix A we present the normalization conventions for the coordinate and momentum eigenstates. In Appendix B we present a detailed derivation of the spectrum of matter operator CU .

2 Review of the 3-string vertex

Let us remind the expressions for the 3-string vertices in the matter and ghost sectors [8, 9] in the presence of the constant background metric $g_{\mu\nu}$:

$$|V_3^{\text{m}}\rangle_{123} = \mathcal{N}_m^{-26} \exp\left[-\frac{1}{2} \sum_{r,s=1}^3 \sum_{m,n=0}^{\infty} (CM^{rs})_{nm} (a_n^{(r)\dagger}, a_m^{(s)\dagger})\right] |\Omega_b\rangle_{123}, \quad (2.1a)$$

$$|V_3^{\text{gh}}\rangle_{123} = \exp\left[-\sum_{r,s=1}^3 \sum_{n=0,m=1}^{\infty} b_{-n}^{(r)} \left(\frac{1}{\sqrt{N}} (C\tilde{M}^{rs}) \sqrt{N}\right)_{nm} c_{-m}^{(s)}\right] |+\rangle_{123}, \quad (2.1b)$$

where

$$CM^{rs} = \frac{1}{3}(C + \alpha^{s-r}U + \alpha^{r-s}\bar{U}), \quad U^\dagger = U, \quad \bar{U} = CUC, \quad U^2 = 1; \quad (2.2a)$$

$$C\tilde{M}^{rs} = \frac{1}{3}(C + \alpha^{s-r}\tilde{U} + \alpha^{r-s}\tilde{\bar{U}}), \quad \tilde{U}^\dagger = \tilde{U}, \quad \tilde{\bar{U}} = C\tilde{U}C, \quad \tilde{U}^2 = 1 \quad (2.2b)$$

and matrices N and C are given by

$$C_{mn} = (-1)^m \delta_{mn} \quad \text{and} \quad N_{nm} = n\delta_{n,m} + \delta_{m,0}\delta_{n,0}. \quad (2.2c)$$

By the bar over the matrix \bar{U} we assume the complex conjugation. The vacuum states are defined as follows

$$a_n|\Omega_b\rangle = 0, \quad n \geq 0; \quad (2.3a)$$

$$b_n|+\rangle = c_n|+\rangle = 0, \quad n \geq 1 \quad \text{and} \quad c_0|+\rangle = 0. \quad (2.3b)$$

The zero mode oscillators $a_{0,\mu}$ and $a_{0,\mu}^\dagger$ are related to the coordinate and momentum modes in the open string expansion as

$$x_\mu = i\frac{\sqrt{\alpha'b}}{2}(a_{0,\mu} - a_{0,\mu}^\dagger) \quad \text{and} \quad p_\mu = \frac{1}{\sqrt{b\alpha'}}(a_{0,\mu} + a_{0,\mu}^\dagger).$$

Here b is a real parameter introduced in [11].

In general the normalization factor \mathcal{N}_m appearing in formula (2.1a) depends on the normalization of the coordinate and momentum eigenstates $|x\rangle$ and $|p\rangle$. We will use the normalization of these states described in Appendix A, for which \mathcal{N}_m is of the form

$$\mathcal{N}_m = \frac{\sqrt{3}}{b} \left(V_{00} + \frac{b}{2} \right) (2\pi\alpha'b)^{\frac{1}{4}}. \quad (2.4)$$

The operators CM and $C\tilde{M}$ are not independent, there is a relation between them. This relation is very much based on the associativity of the Witten's \star -product, and can be obtained from the consideration of 4-string vertices in the matter and ghost sectors. The 4-string vertices in matter and ghost sectors are defined by the matrices V and \tilde{V} correspondingly. As shown in [9] these matrices are related to each other by $\tilde{V} = -\bar{V}$.

The relation between the matrices V and U , which define 4-string and 3-string vertices correspondingly, is given by [8]

$$V + \bar{V} = 4(1 + C(U + \bar{U}))[U + \bar{U} + 4C]^{-1}, \quad (2.5a)$$

$$V - \bar{V} = 2\sqrt{3}(U - \bar{U})C[U + \bar{U} + 4C]^{-1}. \quad (2.5b)$$

The solution of these equations is the following

$$U + \bar{U} = -4(1 - C(V + \bar{V}))[4C - V - \bar{V}]^{-1}, \quad (2.6a)$$

$$U - \bar{U} = -2\sqrt{3}C(V - \bar{V})[4C - V - \bar{V}]^{-1}. \quad (2.6b)$$

These expressions define the function $U = U(V, \bar{V})$. Similar relations to (2.6) can be obtained for the matrices \tilde{U} and $\tilde{\bar{U}}$, one only needs to change $V \mapsto \tilde{V} = -\bar{V}$ [8]. And therefore $\tilde{U} = U(-\bar{V}, -V)^1$ or more precisely

$$\tilde{U} + \tilde{\bar{U}} = -4(1 + C(V + \bar{V}))[4C + V + \bar{V}]^{-1}, \quad (2.7a)$$

$$\tilde{U} - \tilde{\bar{U}} = 2\sqrt{3}C(V - \bar{V})[4C + V + \bar{V}]^{-1}. \quad (2.7b)$$

¹Notice that these relation is only true for $b = 2$.

For the spectroscopy problem it is more convenient to consider matrices CU and UC instead of matrices U and \bar{U} . The reason to do this is that operator CU is a unitary operator. The equations (2.6) and (2.7) allow us to express the operators $C\tilde{U}$ and $\tilde{U}C$ in terms of the operators CU and UC . This relation is of the form

$$C\tilde{U} = -\frac{2CU + 1}{CU + 2} \quad \text{and} \quad \tilde{U}C = -\frac{2UC + 1}{UC + 2}. \quad (2.8)$$

From these relations it follows that eigenvalues of the operators $C\tilde{U}$ and CU are related by the $PSL(2, \mathbb{Z})$ transformation P

$$P(z) = -\frac{2z + 1}{z + 2} = -2 + \frac{3}{z + 2}. \quad (2.9)$$

The transformation P satisfy the equation $(P \circ P)(z) = z$.

3 Spectrum of the operators CU and UC (matter)

3.1 Notations

We have the followings matrices C'_{mn} , C_{ab} , U'_{mn} and U_{ab} , where $a, b \geq 0$ and $m, n \geq 1$. The operators C and C' define BPZ conjugation and are given by $C_{ab} = (-1)^a \delta_{ab}$ and $C'_{mn} = (-1)^m \delta_{mn}$. There operators U and U' have the following properties:

$$U'^{\dagger} = U', \quad \bar{U}' = C'U'C', \quad U'^2 = 1 \quad \text{and} \quad U^{\dagger} = U, \quad \bar{U} = CUC, \quad U^2 = 1, \quad (3.1)$$

where by \bar{U} we denote complex conjugated operator to U . The operators U and U' are related by

$$U'_{mn} = U_{mn} + \frac{1}{1 - U_{00}} U_{m0} U_{0n} \quad \text{and} \quad U_{00} = 1 - \frac{b}{V_{00} + \frac{b}{2}}. \quad (3.2)$$

The vector U_{m0} is an eigenvector of the matrix U' with eigenvalue $+1$:

$$\sum_{n=1}^{\infty} U'_{mn} U_{n0} = U_{m0}, \quad \sum_{m=1}^{\infty} U_{0m} U_{m0} = 1 - U_{00}^2. \quad (3.3)$$

The vectors U_{m0} and U_{0m} are b -dependent, therefore it is convenient to introduce b -independent vectors W_n and \bar{W}_n [11] as

$$W_n = \frac{\sqrt{b}}{1 - U_{00}} U_{0n} \quad \text{and} \quad \bar{W}_n \equiv (-1)^n W_n = \frac{\sqrt{b}}{1 - U_{00}} U_{n0}. \quad (3.4)$$

We also introduce the scalar product (notice that our definition involves complex conjugation):

$$(W, W) \equiv \sum_{n=1}^{\infty} \bar{W}_n W_n \stackrel{3.3}{=} 2V_{00}. \quad (3.5)$$

The expression (3.2) takes the following form in terms of W 's:

$$U'_{mn} = U_{mn} + \frac{\overline{W}_m W_n}{V_{00} + \frac{b}{2}}. \quad (3.6)$$

The matrices $(C'U')_{mn}$ and $(U'C')_{mn}$ have common eigenvectors $v_n^{(\kappa)}$:

$$\sum_{n=1}^{\infty} (C'U')_{mn} v_n^{(\kappa)} = \nu(\kappa) v_m^{(\kappa)} \quad \text{and} \quad \sum_{n=1}^{\infty} (U'C')_{mn} v_n^{(\kappa)} = \overline{\nu}(\kappa) v_m^{(\kappa)}, \quad (3.7)$$

where the components of the vectors $v_m^{(\kappa)}$ are explicitly given by the following generating function [10]

$$f^{(\kappa)}(z) = \sum_{m=1}^{\infty} v_m^{(\kappa)} \frac{z^m}{\sqrt{m}} = \frac{1}{\kappa} (1 - e^{-\kappa \arctan z}). \quad (3.8)$$

Note that since $(C'U')(U'C') = 1$ we get that $|\nu(\kappa)|^2 = 1$. To get an explicit expression for the eigenvalues $\nu(\kappa)$ we will use the eigenvalues $\mu^{rs}(\kappa)$ of the operators M^{rs} [10]

$$\mu^{rs}(\kappa) = \frac{1}{1 + 2 \cosh \frac{\pi\kappa}{2}} \left[1 - 2\delta_{r,s} + e^{\frac{\pi\kappa}{2}} \delta_{r+1,s} + e^{-\frac{\pi\kappa}{2}} \delta_{r,s+1} \right], \quad (3.9a)$$

$$M^{rs} = \frac{1}{3} \left[1 + \alpha^{s-r} C'U' + \alpha^{r-s} U'C' \right]. \quad (3.9b)$$

From the formulae (3.9) one can obtain the following expression for the eigenvalue $\nu(\kappa)$

$$\nu(\kappa) = -\frac{1}{1 + 2 \cosh \frac{\pi\kappa}{2}} \left[2 + \cosh \frac{\pi\kappa}{2} + i\sqrt{3} \sinh \frac{\pi\kappa}{2} \right]. \quad (3.10)$$

The analysis of this formula shows that $\frac{2\pi}{3} < \arg \nu(\kappa) < \frac{4\pi}{3}$ (see Figure 1).

The eigenvectors $v^{(\kappa)}$ belong to the continuous spectrum. As it follows from the expression (3.10) the spectrum of operators $C'U'$ and $U'C'$ is non-degenerate.

The normalization of the vectors $v^{(\kappa)}$ is given by [12]

$$(v^{(\kappa)}, v^{(\kappa')}) = \mathcal{N}(\kappa) \delta(\kappa - \kappa'), \quad \text{where} \quad \mathcal{N}(\kappa) = \frac{2}{\kappa} \sinh \frac{\pi\kappa}{2}. \quad (3.11)$$

Since the operator M^{11} (3.9b) is a hermitian operator and $v^{(\kappa)}$ are its only eigenvectors [10] we conclude that the vectors $v^{(\kappa)}$ form a complete set:

$$\delta_{mn} = \int d\kappa \frac{1}{\mathcal{N}(\kappa)} v_m^{(\kappa)} v_n^{(\kappa)}. \quad (3.12)$$




Figure 1: On the picture we have presented the spectrum of the operator CU . It consist of continuous spectrum, which is represented by the function $\nu(\kappa)$ (solid blue line), and discrete spectrum, which contains two points ξ and $\bar{\xi}$ (red dots).

The scalar products among W_n and $v_n^{(\kappa)}$ can be obtained from the paper² [13] and are given by

$$(v^{(\kappa)}, W) = -\frac{\sqrt{2}}{\kappa} \frac{\cosh \frac{\pi\kappa}{2} - 1 - i\sqrt{3} \sinh \frac{\pi\kappa}{2}}{2 \cosh \frac{\pi\kappa}{2} + 1} \equiv -\frac{\sqrt{2}}{\kappa} [1 + \nu(\kappa)]. \quad (3.13a)$$

$$(v^{(\kappa)}, \bar{W}) = -\frac{\sqrt{2}}{\kappa} \frac{\cosh \frac{\pi\kappa}{2} - 1 + i\sqrt{3} \sinh \frac{\pi\kappa}{2}}{2 \cosh \frac{\pi\kappa}{2} + 1} \equiv -\frac{\sqrt{2}}{\kappa} [1 + \bar{\nu}(\kappa)], \quad (3.13b)$$

3.2 Eigenvectors of the operators CU and UC .

We want to find a spectrum of the operators CU and UC . Since $UC = (CU)^\dagger$ it is enough to find the spectrum of the operator CU . To this end we have to solve the equation

$$\sum_{b=0}^{\infty} (CU)_{ab} V_b^{(\xi)} = \xi V_a^{(\xi)}. \quad (3.14)$$

Notice that since CU is a unitary operator all of its eigenvalues ξ belong to the unit circle. Consideration of the two cases $a \neq 0$ and $a = 0$ yields the equations:

$$\sum_{n=1}^{\infty} (CU)_{mn} V_n^{(\xi)} + \sum_{n=1}^{\infty} C_{mn} U_{n0} V_0^{(\xi)} = \xi V_m^{(\xi)}, \quad (3.15a)$$

$$\sum_{n=1}^{\infty} U_{0n} V_n^{(\xi)} + U_{00} V_0^{(\xi)} = \xi V_0^{(\xi)}. \quad (3.15b)$$

²To this end one have to use expression (3.11) from [13] and the following relation between v_e , v_o and W :

$$W = -\sqrt{2}(v_e - iv_o) \quad \text{and} \quad \bar{W} = -\sqrt{2}(v_e + iv_o).$$

Notice that this relation is different from the one presented in [13]. The origin of these difference is explained in the end of Appendix A of [11].

Using the expression (3.6) one (3.4) one can rewrite the equation (3.15a) in the following form:

$$\sum_{n=1}^{\infty} \left[(C'U')_{mn} - \xi \delta_{mn} \right] V_n^{(\xi)} = \frac{1}{\sqrt{b}} W_m \sum_{n=1}^{\infty} U_{0n} V_n^{(\xi)} - \frac{1 - U_{00}}{\sqrt{b}} W_m V_0^{(\xi)}.$$

Further simplification can be achieved by use of equation (3.15b). Finally the equations we have to solve get the following simple form

$$\sum_{n=1}^{\infty} \left[(C'U')_{mn} - \xi \delta_{mn} \right] V_n^{(\xi)} = W_m \frac{V_0^{(\xi)}}{\sqrt{b}} (\xi - 1), \quad (3.16a)$$

$$(\overline{W}, V^{(\xi)}) = \frac{V_0^{(\xi)}}{\sqrt{b}} \left[\left(V_{00} + \frac{b}{2} \right) \xi - \left(V_{00} - \frac{b}{2} \right) \right]. \quad (3.16b)$$

To solve these equations it is very important to notice that the operator $C'U'$ has only continuous spectrum. In this case one can easily write the solution to the equation (3.16a). We have to distinguish two different cases:

1. The number ξ is in the spectrum of $C'U'$. This means that there is a real number κ such that $\xi = \nu(\kappa)$. In this case the solution of the (3.16a) is given by

$$V_m^{(\kappa)} = A^{(\kappa)} v_m^{(\kappa)} - [1 - \nu(\kappa)] \frac{V_0^{(\kappa)}}{\sqrt{b}} \left(\mathcal{P} \frac{1}{C'U' - \nu(\kappa)} \right)_{mn} W_n, \quad (3.17a)$$

where $A^{(\kappa)}$ is an arbitrary number and $\mathcal{P} \frac{1}{x}$ means principal value. The only reason why we have chosen the principal value instead of for example $\frac{1}{x+i0}$ is that the formulae will be a bit simpler for this case. Substitution of this solution into (3.16b) allows us to determine $A^{(\kappa)}$ in terms of $V_0^{(\kappa)}$:

$$A^{(\kappa)} = \frac{V_0^{(\kappa)}}{\sqrt{b}} \frac{\nu(\kappa) - 1}{(\overline{W}, v^{(\kappa)})} \left[V_{00} + \frac{b}{2} \frac{\nu(\kappa) + 1}{\nu(\kappa) - 1} - (\overline{W}, \mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W) \right] \quad (3.17b)$$

As we will see equations (3.17) define a continuous spectrum of the operator CU . This continuous spectrum is *non-degenerate* and consists of the vectors $V^{(\kappa)}$ with eigenvalues $\nu(\kappa)$.

2. If ξ is not in the spectrum of $C'U'$, the solution of the equation (3.16a) is given by

$$V_m^{(\xi)} = \frac{V_0^{(\xi)}}{\sqrt{b}} (\xi - 1) \left(\frac{1}{C'U' - \xi} \right)_{mn} W_n. \quad (3.18a)$$

Substitution of this solution into (3.16b) yields the equation on the eigenvalue ξ (discrete spectrum):

$$(\overline{W}, \frac{1}{C'U' - \xi} W) = V_{00} + \frac{b}{2} \frac{\xi + 1}{\xi - 1}. \quad (3.18b)$$

This equation in general has two complex conjugated solutions belonging to the unit circle (see Figure 1). One can easily check that the vector $CV^{(\xi)}$ is also an eigenvector of the operator CU with the eigenvalue $\bar{\xi}$.

The details of solving of these equations are presented in Appendix B. The results are collected in the next subsection.

3.3 Summary

The spectrum of the operators CU and UC consist of two branches: continuous and discrete.

- **Discrete spectrum**

The discrete spectrum of the operator CU consist of two points ξ and $\bar{\xi}$ (see Figure 1). They are given as the solutions of the equation

$$\Re F(\eta) = \frac{b}{4}, \quad (3.19)$$

where

$$F(\eta) = \psi\left(\frac{1}{2} + \frac{\eta}{2\pi i}\right) - \psi\left(\frac{1}{2}\right)$$

and ξ is related to η by

$$\xi = -\frac{1}{1 - 2 \cosh \eta} \left[2 - \cosh \eta - i\sqrt{3} \sinh \eta \right]. \quad (3.20)$$

For $b > 0$ equation (3.19) has two real solutions. On the Figure 4 we presented the $\arg \bar{\xi}(b)$ for this solution.

The generating functions for the eigenvectors from the discrete spectrum is given by

$$F^{(\xi)}(z) \equiv \sum_{n=1}^{\infty} V_n^{(\xi)} \frac{z^n}{\sqrt{n}} = -\sqrt{\frac{2}{b}} V_0^{(\xi)} \left[\frac{b}{4} + \frac{\pi}{2\sqrt{3}} \frac{\xi - 1}{\xi + 1} + \log iz \right. \\ \left. + e^{-2i(1+\frac{\eta}{\pi i}) \arctan z} \text{LerchPhi}\left(e^{-4i \arctan z}, 1, \frac{1}{2} + \frac{\eta}{2\pi i}\right) \right], \quad (3.21a)$$

where

$$x \text{LerchPhi}(w, 1, x) = {}_2F_1(1, x; x + 1; w).$$

The action of the operator C on the eigenvectors can be read from the formulae

$$F^{(\bar{\xi})}(z) \equiv F^{(\xi)}(-z) \iff CV^{(\xi)} = V^{(\bar{\xi})}. \quad (3.21b)$$

The norm of the vectors from the discrete spectrum is given by

$$((V^{(\xi)}, V^{(\xi)})) = |V_0^{(\xi)}|^2 \sinh \eta \frac{\partial}{\partial \eta} \left[\log \Re F(\eta) \right]. \quad (3.21c)$$

Further we will assume that $V_0^{(\xi)}$ is chosen in such way (see (B.17)) that the norm (3.21c) is equal to one.

- **Continuous spectrum**

The continuous spectrum of the operator CU is given by the function $\nu(\kappa)$, $\kappa \in \mathbb{R}$ (see Figure 1)

$$\nu(\kappa) = -\frac{1}{1 + 2 \cosh \frac{\pi\kappa}{2}} \left[2 + \cosh \frac{\pi\kappa}{2} + i\sqrt{3} \sinh \frac{\pi\kappa}{2} \right]. \quad (3.22)$$

The generating function for the eigenvector corresponding to eigenvalue $\nu(\kappa)$ is given by

$$\begin{aligned} F^{(\kappa)}(z) \equiv \sum_{n=1}^{\infty} V_n^{(\kappa)} \frac{z^n}{\sqrt{n}} = & V_0^{(\kappa)} \left[\frac{2}{b} \right]^{\frac{1}{2}} \left[-\frac{b}{4} - \left(\Re F_c(\kappa) - \frac{b}{4} \right) e^{-\kappa \arctan z} \right. \\ & - \left(\frac{\pi}{2\sqrt{3}} \frac{\nu - 1}{\nu + 1} + \frac{2i}{\kappa} \right) - \log iz + 2if^{(\kappa)}(z) \\ & \left. - \text{LerchPhi}(e^{-4i \arctan z}, 1, 1 + \frac{\kappa}{4i}) e^{-4i \arctan z} e^{-\kappa \arctan z} \right]. \end{aligned} \quad (3.23a)$$

The action of the operator C' on the eigenvectors can be obtained from the identity on generating functions

$$F^{(\kappa)}(-z) = F^{(-\kappa)}(z) \iff CV^{(\kappa)} = V^{(-\kappa)}. \quad (3.23b)$$

The norm of the vectors from continuous spectrum is given by

$$((V^{(\kappa)}, V^{(\kappa')})) = \frac{2}{b} |V_0^{(\kappa)}|^2 \mathcal{N}(\kappa) \left[4 + \kappa^2 \left(\Re F_c(\kappa) - \frac{b}{4} \right)^2 \right] \delta(\kappa - \kappa'). \quad (3.23c)$$

Further we will assume that $V_0^{(\kappa)}$ is real and chosen in such way (see (B.33)) that the vectors from continuous spectrum (3.23c) are normalized to δ -function without any additional factors.

Partition of the unity.

Since the operator $CM = \frac{1}{3}(1 + CU + UC)$ is a hermitian operator and vectors $V^{(\kappa)}$, $V^{(\xi)}$ and $V^{(\bar{\xi})}$ are the only its eigenvectors we conclude that these vectors form a complete set. Therefore we can write the following partition of the unit operator Id

$$\text{Id} = \int_{-\infty}^{\infty} d\kappa V^{(\kappa)} \otimes V^{(\kappa)} + V^{(\xi)} \otimes V^{(\xi)} + V^{(\bar{\xi})} \otimes V^{(\bar{\xi})}, \quad (3.24)$$

where we assume that all vectors are normalized to 1, i.e. their zero components have been chosen as in equations (B.33) and (B.17).




Figure 2: On the pane a) we show the spectrum of the operator CU , which defines the \star -product in the matter sector. On the pane b) we show the spectrum of the operator $C\tilde{U}$, which defines the \star -product in the ghost sector. The arrow between the panes shows the relation among these two operators.

4 Spectrum of the operators defining ghost vertex

4.1 Spectrum of the operators $C\tilde{U}$ and $\tilde{U}C$ (ghosts)

In the section 2 we have shown that the operator $C\tilde{U}$ for ghost sector and the operator CU for the matter sector are related by $PSL(2, \mathbb{Z})$ transformation P (2.9). From this it follows that the eigenvectors of the operator $C\tilde{U}$ are the same as of those CU , and the eigenvalues are related by the transformation P . The resulting spectrum of the operator $C\tilde{U}$ is presented on Figure 2b).

So the spectrum of $C\tilde{U}$ consists of two branches: continuous and discrete.

- **Discrete spectrum**

The discrete spectrum of operator $C\tilde{U}$ consist of two points $\tilde{\xi}$ and $\bar{\tilde{\xi}}$, which are defined as

$$\tilde{\xi} \equiv P(\xi) = -\frac{2 + \cosh \eta - i\sqrt{3} \sinh \eta}{1 + 2 \cosh \eta}. \quad (4.1)$$

The eigenvectors corresponding to these eigenvalues are $V^{(\tilde{\xi})} \equiv V^{(\xi)}$ and $V^{(\bar{\tilde{\xi}})} \equiv V^{(\bar{\xi})}$, and their generating functions are given by equation (3.21a).

- **Continuous spectrum** The discrete spectrum of operator $C\tilde{U}$ consists of an arc shown on the Figure 2. The parametrization of this arc is given by function $\tilde{\nu}(\kappa)$

$$\tilde{\nu}(\kappa) \equiv P(\nu(\kappa)) = -\frac{2 - \cosh \frac{\pi\kappa}{2} + i\sqrt{3} \sinh \frac{\pi\kappa}{2}}{1 - 2 \cosh \frac{\pi\kappa}{2}}. \quad (4.2)$$

The eigenvectors corresponding to these eigenvalues are $V^{(\kappa)}$, and their generating functions are given by equation (3.23a).

Partition of the unity.

Partition of the unit operator is the same as in (3.24).

4.2 Spectrum of the operators $C'\tilde{U}'$ and $\tilde{U}'C'$ (ghosts)

By analogy to the matter sector the operators $C\tilde{U}$ and $\tilde{U}C$ can be presented in the following block from

$$C\tilde{U} = \begin{pmatrix} \tilde{U}_{00} & K^t \\ K & C'\tilde{U}' + \lambda K \otimes K^t \end{pmatrix}, \quad (4.3)$$

where λ is fixed by the condition that $C\tilde{U}$ is unitary operator, i.e.

$$\lambda = \frac{1}{\tilde{U}_{00} \pm 1}, \quad \tilde{U}'\bar{K} = \mp \bar{K} \quad \text{and} \quad \sum_{n=1}^{\infty} \bar{K}_n K_n = 1 - \tilde{U}_{00}^2. \quad (4.4)$$

Using the fact that operators $C\tilde{U}$ and CU (for $b=2$) are related by the transformation (2.9) we can easily obtain the expressions for \tilde{U}_{00} , λ , K and $C'\tilde{U}'$

$$\tilde{U}_{00} = \frac{1 - \tilde{V}_{00}}{1 + \tilde{V}_{00}}, \quad \tilde{V}_{00} = 3 \log 3 \quad \lambda = (1 + \tilde{U}_{00})^{-1}; \quad (4.5a)$$

$$K = -\frac{3}{\sqrt{2}}(\tilde{U}_{00} + 1)(C'U' + 2)^{-1}W, \quad C'\tilde{U}' = P(C'U'). \quad (4.5b)$$

Now we can conclude that the spectrum of the operator $C'\tilde{U}'$ is only continuous and consists of the eigenvectors defined by generating function (3.8) and corresponding eigenvalues are given by equation (4.2).

Basically only the right column of the operator (4.3) is appeared in the ghost vertex (2.1b). Therefore we need to know how represent this column in the diagonal form. In principal it is possible to diagonalize the matrix $(CU)_{mn}$, $m, n \geq 1$. To this end notice that the matrix $(CU)_{mn}$ is not a unitary one, i.e.

$$\sum_{n=1}^{\infty} (UC)_{mn}(CU)_{nk} = \delta_{nk} - \bar{K}_n \otimes K_k.$$

The spectrum of this operator consists of two branches continuous and discrete. The eigenvalues from the continuous spectrum are represented by the function $\tilde{\nu}(\kappa)$, and the corresponding eigenvectors are complex (this is very different from the cases we had before). There are two points in the discrete spectrum, one eigenvalue is inside the unite circle and another one is outside. The corresponding eigenvectors are also complex. Since all the eigenvectors are complex it looks impossible to apply them to diagonalize the vertex (4.3).

So will diagonalize only the operator $C'\tilde{U}'$ and express the vector K in terms of $v^{(\kappa)}$. The scalar product of K and \bar{K} with $v^{(\kappa)}$ is given by

$$(v^{(\kappa)}, K) = (1 + \tilde{U}_{00}) \frac{1 - \tilde{\nu}(\kappa)}{\kappa} \quad \text{and} \quad (v^{(\kappa)}, \bar{K}) = (1 + \tilde{U}_{00}) \frac{1 - \tilde{\bar{\nu}}(\kappa)}{\kappa}. \quad (4.6)$$

5 Diagonal representation of the open string star

In this section we will rewrite the 3-string vertex in the matter and ghost sectors in the oscillator basis related to the eigenvectors of operators CU and UC ($C\tilde{U}$ and $\tilde{U}C$) found in the previous section.

5.1 Matter 3-vertex

Let us introduce new oscillators

$$a_\kappa = \sum_{n=0}^{\infty} V_n^{(\kappa)} a_n, \quad a_\xi = \sum_{n=0}^{\infty} V_n^{(\xi)} a_n \quad \text{and} \quad a_{\bar{\xi}} = \sum_{n=0}^{\infty} V_n^{(\bar{\xi})} a_n. \quad (5.1a)$$

Using the completeness of the basis (3.24) we can express old oscillators as

$$a_n = \int_{-\infty}^{\infty} d\kappa V_n^{(\kappa)} a_\kappa + V_n^{(\xi)} a_\xi + V_n^{(\bar{\xi})} a_{\bar{\xi}}. \quad (5.1b)$$

The commutation relations for oscillators a_κ are of the form

$$[a_{\kappa,\mu}, a_{\kappa',\nu}^\dagger] = g_{\mu\nu} \delta(\kappa - \kappa') \quad \text{and} \quad [a_{\kappa,\mu}, a_{\kappa',\nu}] = 0. \quad (5.1c)$$

The operator C acts on the on the new oscillators in the following way

$$Ca_\kappa = a_{-\kappa} \quad \text{and} \quad Ca_\xi = a_{\bar{\xi}}. \quad (5.1d)$$

Substitution of the expression (5.1b) into (2.1a) yields the following representation for the matter 3-string vertex

$$|V_3^m\rangle_{123} = \mathcal{N}_m^{-26} \exp \left[-\frac{1}{2} \sum_{r,s} \int_{-\infty}^{\infty} d\kappa \mu^{rs}(\kappa) (Ca_{\kappa}^{(r)\dagger}, a_{\kappa}^{(s)\dagger}) - \sum_{r,s} \mu_{\xi}^{rs} (Ca_{\xi}^{(r)\dagger}, a_{\xi}^{(s)\dagger}) \right] |\Omega_b\rangle_{123}, \quad (5.2)$$

where

$$\mu^{rs}(\kappa) = \frac{1}{3} [1 + \alpha^{s-r} \nu(\kappa) + \alpha^{r-s} \bar{\nu}(\kappa)] = \frac{1 - 2\delta_{r,s} + e^{\frac{\pi\kappa}{2}} \delta_{r+1,s} + e^{-\frac{\pi\kappa}{2}} \delta_{r,s+1}}{1 + 2 \cosh \frac{\pi\kappa}{2}}, \quad (5.3a)$$

$$\mu_{\xi}^{rs} = \frac{1}{3} [1 + \alpha^{s-r} \xi + \alpha^{r-s} \bar{\xi}] = \frac{1 - 2\delta_{r,s} - e^{\eta} \delta_{r+1,s} - e^{-\eta} \delta_{r,s+1}}{1 - 2 \cosh \eta}. \quad (5.3b)$$

Equation (5.2) gives almost diagonal representation of the interaction vertex. To write it in diagonal form we need to introduce oscillator basis which has definite parity with respect to the operator C . To this end let us introduce even and odd oscillator modes

$$e_\kappa = \frac{a_\kappa + Ca_\kappa}{\sqrt{2}}, \quad o_\kappa = \frac{a_\kappa - Ca_\kappa}{i\sqrt{2}}, \quad e_\xi = \frac{a_\xi + Ca_\xi}{\sqrt{2}}, \quad o_\xi = \frac{a_\xi - Ca_\xi}{i\sqrt{2}}; \quad (5.4a)$$

$$a_\kappa = \frac{e_\kappa + io_\kappa}{\sqrt{2}}, \quad Ca_\kappa = \frac{e_\kappa - io_\kappa}{\sqrt{2}}, \quad a_\xi = \frac{e_\xi + io_\xi}{\sqrt{2}}, \quad Ca_\xi = \frac{e_\xi - io_\xi}{\sqrt{2}}. \quad (5.4b)$$

We have introduced a factor i in the definition of the odd oscillators so that along with the even ones they satisfy the same BPZ conjugation property [6]

$$\text{bpz } o_\kappa = -o_\kappa^\dagger \quad \text{and} \quad \text{bpz } e_\kappa = -e_\kappa^\dagger. \quad (5.5)$$

Substitution of (5.4b) into the expression for 3-string vertex (5.2) yields

$$\begin{aligned} |V_3\rangle = \mathcal{N}_m^{-26} \exp & \left[-\frac{1}{4} \sum_{r,s} \int_0^\infty d\kappa (\mu^{rs} + \mu^{sr}) \left\{ (e_\kappa^{(r)\dagger}, e_\kappa^{(s)\dagger}) + (o_\kappa^{(r)\dagger}, o_\kappa^{(s)\dagger}) \right\} \right. \\ & - \frac{i}{4} \sum_{r,s} \int_0^\infty d\kappa (\mu^{rs} - \mu^{sr}) \left\{ (e_\kappa^{(r)\dagger}, o_\kappa^{(s)\dagger}) - (o_\kappa^{(r)\dagger}, e_\kappa^{(s)\dagger}) \right\} \\ & - \frac{1}{4} \sum_{r,s} (\mu_\xi^{rs} + \mu_\xi^{sr}) \left\{ (e_\xi^{(r)\dagger}, e_\xi^{(s)\dagger}) + (o_\xi^{(r)\dagger}, o_\xi^{(s)\dagger}) \right\} \\ & \left. - \frac{i}{4} \sum_{r,s} (\mu_\xi^{rs} - \mu_\xi^{sr}) \left\{ (e_\xi^{(r)\dagger}, o_\xi^{(s)\dagger}) - (o_\xi^{(r)\dagger}, e_\xi^{(s)\dagger}) \right\} \right] |\Omega_b\rangle_{123}. \quad (5.6) \end{aligned}$$

So this expression gives the diagonal representation of the full 3-string interaction vertex in the matter sector. Using this representation in the next section we will show that it encodes a family of even Moyal algebras [6].

Notice that for $\kappa = 0$ the odd oscillator $o_{\kappa=0}$ is equal to zero and we are left only with the even oscillator $e_{\kappa=0}$ (see (B.26)). To derive the meaning of this mode it is convenient to introduce coordinate and momentum oscillators

$$x_\kappa = \frac{i}{\sqrt{2}}(a_\kappa - a_\kappa^\dagger) \quad \text{and} \quad p_\kappa = \frac{1}{\sqrt{2}}(a_\kappa + a_\kappa^\dagger).$$

Using formulae (B.26) and (B.33) we can write $x_{\kappa=0}$ and $p_{\kappa=0}$ in the following way

$$x_{\kappa=0} = \frac{1}{2\sqrt{\pi\alpha'}} \left[x_0 + 2\sqrt{\alpha'} \sum_{n=1}^{\infty} (-1)^n x_{2n} \right] = \frac{1}{2\sqrt{\pi\alpha'}} X\left(\frac{\pi}{2}\right); \quad (5.7a)$$

$$p_{\kappa=0} = \sqrt{\pi\alpha'} \left[\frac{1}{2\pi} p_0 + \frac{1}{\pi\sqrt{\alpha'}} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n} p_{2n} \right]. \quad (5.7b)$$

So we get that one of the variables is proportional to $X(\frac{\pi}{2})$.

5.2 Ghost 3-vertex

Let us first rewrite the ghost vertex (2.1b) using the representation (4.3) for the ghost operators CU . To this end we notice that

$$(C\widetilde{M}^{rs})_{0m} = \frac{1}{3}\alpha^{s-r} K_m + \frac{1}{3}\alpha^{r-s} \overline{K}_m, \quad (5.8a)$$

$$(C\widetilde{M}^{rs})_{nm} = (C\widetilde{M}'^{rs})_{nm} + \frac{1}{3(1+\widetilde{U}_{00})} \left[\alpha^{s-r} \overline{K}_n \otimes K_m + \alpha^{r-s} K_n \otimes \overline{K}_m \right]. \quad (5.8b)$$

Substitution of these expressions into (2.1b) yields

$$\begin{aligned}
|V_3^{\text{gh}}\rangle_{123} = \exp & \left[-\frac{1}{3} \sum_{r,s=1}^3 \sum_{m=1}^{\infty} b_0^{(r)} \left[\alpha^{s-r} K_m + \frac{1}{3} \alpha^{r-s} \bar{K}_m \right] \sqrt{m} c_{-m}^{(s)} \right. \\
& - \sum_{r,s=1}^3 \sum_{m,n=1}^{\infty} b_{-n}^{(r)} \frac{1}{\sqrt{n}} (C \widetilde{M}'_{rs})_{nm} \sqrt{m} c_{-m}^{(s)} \\
& \left. - \frac{1}{3(1 + \tilde{U}_{00})} \sum_{r,s=1}^3 \sum_{m,n=1}^{\infty} b_{-n}^{(r)} \frac{1}{\sqrt{n}} \left[\alpha^{s-r} \bar{K}_n \otimes K_m + \alpha^{r-s} K_n \otimes \bar{K}_m \right] \sqrt{m} c_{-m}^{(s)} \right] |+\rangle_{123}, \quad (5.9)
\end{aligned}$$

Now it is obvious that we have to introduce the following oscillators

$$b_{\kappa}^{\dagger} = \frac{1}{\sqrt{\mathcal{N}(\kappa)}} \sum_{n=1}^{\infty} \frac{v_n^{(\kappa)}}{\sqrt{n}} b_{-n}, \quad b^{\dagger} = \frac{1}{\sqrt{1-\tilde{U}_{00}^2}} \sum_{n=1}^{\infty} \frac{K_n}{\sqrt{n}} b_{-n}^{\dagger}, \quad \bar{b}^{\dagger} = \frac{1}{\sqrt{1-\tilde{U}_{00}^2}} \sum_{n=1}^{\infty} \frac{\bar{K}_n}{\sqrt{n}} b_{-n}; \quad (5.10a)$$

$$c_{\kappa}^{\dagger} = \frac{1}{\sqrt{\mathcal{N}(\kappa)}} \sum_{n=1}^{\infty} v_n^{(\kappa)} \sqrt{n} c_{-n}, \quad c^{\dagger} = \frac{1}{\sqrt{1-\tilde{U}_{00}^2}} \sum_{n=1}^{\infty} K_n \sqrt{n} c_{-n}, \quad \bar{c}^{\dagger} = \frac{1}{\sqrt{1-\tilde{U}_{00}^2}} \sum_{n=1}^{\infty} \bar{K}_n \sqrt{n} c_{-n}. \quad (5.10b)$$

Using formulae (4.6) we can easily express the oscillators c^{\dagger} and \bar{c}^{\dagger} through c_{κ}^{\dagger} (and the same for b^{\dagger} and \bar{b}^{\dagger})

$$c^{\dagger} = \int_{-\infty}^{\infty} d\kappa \lambda(\kappa) c_{\kappa}^{\dagger} \quad \text{and} \quad \bar{c}^{\dagger} = \int_{-\infty}^{\infty} d\kappa \bar{\lambda}(\kappa) c_{\kappa}^{\dagger}, \quad (5.11a)$$

$$\text{where} \quad \lambda(\kappa) = [\tilde{V}_{00} \mathcal{N}(\kappa)]^{-\frac{1}{2}} \frac{1 - \tilde{\nu}(\kappa)}{\kappa}. \quad (5.11b)$$

Using the completeness (3.12) of the basis $v^{(\kappa)}$ we can express old oscillators as

$$c_{-n} = \int_{-\infty}^{\infty} d\kappa v_n^{(\kappa)} c_{\kappa}^{\dagger} \quad \text{and} \quad b_{-n} = \int_{-\infty}^{\infty} d\kappa v_n^{(\kappa)} b_{\kappa}^{\dagger}. \quad (5.12)$$

The commutation relations for oscillators b_{κ} , c_{κ} , b and c are of the form

$$\{b_{\kappa}, c_{\kappa'}^{\dagger}\} = \delta(\kappa - \kappa'), \quad \{b_{\kappa}, c^{\dagger}\} = \lambda(\kappa) \quad \text{and} \quad \{b_{\kappa}, \bar{c}^{\dagger}\} = \bar{\lambda}(\kappa). \quad (5.13)$$

The operator C acts on the on the new oscillators in the following way

$$C c_{\kappa}^{\dagger} = -c_{-\kappa}^{\dagger} \quad \text{and} \quad C c^{\dagger} = \bar{c}^{\dagger} \quad (5.14)$$

and the same for b_κ , b and \bar{b} . Substitution of the expression (5.12) into (2.1b) yields the following representation for the matter 3-string vertex

$$|V_3^{\text{gh}}\rangle_{123} = \exp \left[-\frac{1}{3}(1 - \tilde{U}_{00}^2)^{\frac{1}{2}} \sum_{r,s} b_0^{(r)\dagger} (\alpha^{s-r} c^{(s)\dagger} + \alpha^{r-s} \bar{c}^\dagger) \right. \\ \left. + \sum_{r,s} \int_0^\infty d\kappa \tilde{\mu}^{rs}(\kappa) \left[b_\kappa^{(r)\dagger} c_{-\kappa}^{(s)\dagger} + b_\kappa^{(s)\dagger} c_{-\kappa}^{(r)\dagger} \right] \right. \\ \left. - \frac{1 - \tilde{U}_{00}}{3} \sum_{r,s} \left(\alpha^{s-r} \bar{b}^{(r)\dagger} c^{(s)\dagger} + \alpha^{r-s} b^{(r)\dagger} \bar{c}^{(s)\dagger} \right) \right] |+\rangle_{123}, \quad (5.15)$$

where

$$\tilde{\mu}^{rs}(\kappa) = \frac{1}{3} \left[1 + \alpha^{s-r} \tilde{\nu}(\kappa) + \alpha^{r-s} \bar{\nu}(\kappa) \right] = \frac{-1 + 2\delta_{r,s} - e^{\frac{\pi\kappa}{2}} \delta_{r+1,s} - e^{-\frac{\pi\kappa}{2}} \delta_{r,s+1}}{1 - 2 \cosh \frac{\pi\kappa}{2}}, \quad (5.16)$$

The equation (5.15) gives almost diagonal representation of the interaction vertex. To write it in the diagonal form one can introduce oscillator basis which has definite parity with respect to the operator C .

6 Moyal structure

The aim of this section is to identify Moyal algebras appeared in the expression for the 3-string vertex in the matter sector.

6.1 Oscillator vertex for the Moyal product

Let us remind the expression for the vertex representation of the Moyal product obtained in [6]. To this end we have to introduce the Moyal product on functions and the function-state correspondence.

For our purpose it is sufficient to consider functions on the two-dimensional real space \mathbb{R}^2 . The Moyal product is given by the following integral operator

$$(f * g)(x_1) = \int_{\mathbb{R}^2 \times \mathbb{R}^2} d^2x_2 d^2x_3 K(x_1, x_2, x_3) f(x_2) g(x_3), \quad (6.1a)$$

where the kernel $K(x_1, x_2, x_3)$ is given by

$$K(x_1, x_2, x_3) = \frac{1}{\pi^2 \det \Theta} \exp \left(-2i \left[x_1^\alpha (\Theta^{-1})_{\alpha\beta} x_2^\beta + x_2^\alpha (\Theta^{-1})_{\alpha\beta} x_3^\beta + x_3^\alpha (\Theta^{-1})_{\alpha\beta} x_1^\beta \right] \right), \quad (6.1b)$$

indexes $\alpha, \beta = 1, 2$ and we choose the coordinate system in which the antisymmetric tensor $\Theta^{\alpha\beta}$ has the diagonal form

$$\Theta^{\alpha\beta} = \begin{pmatrix} 0 & \theta \\ -\theta & 0 \end{pmatrix}. \quad (6.1c)$$

The function state correspondence is defined as

$$f \leftrightarrow |f\rangle = \int d^2x f(x) |x\rangle \quad (6.2a)$$

and conjugated state is given by

$$\langle f| = \int d^2x \langle x| f(x), \quad (6.2b)$$

where $|x\rangle$ are eigenvectors of the coordinate operator and they normalization is given by (A.4a). To be able to introduce the vertex operator we need to rewrite the state $|x\rangle$ in terms of creation/annihilation operators. The creation/annihilation operators a^\dagger, b^\dagger and a, b are related to coordinate x^α and momentum p_α operators by formulae

$$x_1 = \frac{i}{\sqrt{2}}(a - a^\dagger), \quad p_1 = \frac{1}{\sqrt{2}}(a + a^\dagger) \quad \text{and} \quad x_2 = \frac{i}{\sqrt{2}}(b - b^\dagger), \quad p_2 = \frac{1}{\sqrt{2}}(b + b^\dagger). \quad (6.3a)$$

The state $\langle x|$ can be written through the operators $a_\alpha, a_\alpha^\dagger$ (we assume $a_\alpha = (a, b)$) in the following form

$$\langle x| = \frac{1}{\sqrt{2\pi}} \langle 0| \exp\left[-\frac{1}{2}(x, x) - i\sqrt{2}(a, x) + \frac{1}{2}(a, a)\right]. \quad (6.3b)$$

Expressions for the conjugated state and eigenstates of momentum operator are presented in Appendix A (for $b\alpha' = 2, d = 2$).

Now we can write the Moyal product in terms of vertices

$$|f * g\rangle_1 = ({}_2\langle f| \otimes {}_3\langle g|) |V_3(\theta)\rangle_{123}, \quad (6.4)$$

where the vertex $|V_3(\theta)\rangle$ has the following form³ [6]

$$\begin{aligned} |V_3(\theta)\rangle_{123} = & \frac{2}{3\sqrt{\pi}} \frac{1}{1 + \frac{\theta^2}{12}} \exp\left[-\frac{1}{2} \frac{\theta^2 - 4}{\theta^2 + 12} (a^{(1)\dagger} a^{(1)\dagger} + b^{(1)\dagger} b^{(1)\dagger} + c.p.) \right. \\ & - \frac{8}{\theta^2 + 12} (a^{(1)\dagger} a^{(2)\dagger} + b^{(1)\dagger} b^{(2)\dagger} + c.p.) \\ & \left. - \frac{4i\theta}{\theta^2 + 12} (a^{(1)\dagger} b^{(2)\dagger} - b^{(1)\dagger} a^{(2)\dagger} + c.p.) \right] |0\rangle_{123}. \quad (6.5) \end{aligned}$$

Notice that we use a bit different normalization of coordinate and momentum eigenstates as compared to [6], but the final form of the vertex (6.5) is precisely the same.

³ The other way to obtain this expression is to notice that in the momentum representation the 3-vertex corresponding to the Moyal product is given by

$$\begin{aligned} |V_3\rangle_{123} = & \frac{1}{2\pi} \int dp^{(1)} dp^{(2)} dp^{(3)} \delta^{(2)}(p^{(1)} + p^{(2)} + p^{(3)}) \\ & \times \exp\left[\frac{i}{6} \Theta^{\alpha\beta} (p_\alpha^{(1)} p_\beta^{(2)} + p_\alpha^{(2)} p_\beta^{(3)} + p_\alpha^{(3)} p_\beta^{(1)})\right] |p^{(1)}\rangle_1 \otimes |p^{(2)}\rangle_2 \otimes |p^{(3)}\rangle_3. \end{aligned}$$

To obtain the expression (6.5) one needs to substitute states $|p\rangle$ from (A.1) and integrate over momenta.

6.2 Identification of Moyal structures

In this subsection we construct a correspondence of the open string \star -product to the Moyal product. Basically we will single out two types of Moyal algebra (continuous and discrete) in the expression (5.6) for the 3-string vertex. We show that deformation parameter of the continuous Moyal algebra is parametrized by $\kappa \in [0, \infty)$ and the one of discrete algebra is related to η .

Let us now try to identify the vertex (6.5), which represent the Moyal product, in the diagonal matter 3-string vertex (5.6). To this end we need to identify

$$\{a_\mu^\dagger, b_\mu^\dagger\} \leftrightarrow \{e_{\kappa,\mu}^\dagger, o_{\kappa,\mu}^\dagger\} \quad \text{or} \quad \{a_\mu^\dagger, b_\mu^\dagger\} \leftrightarrow \{e_{\xi,\mu}^\dagger, o_{\xi,\mu}^\dagger\}. \quad (6.6)$$

Here we add indexes μ to the the oscillators a^\dagger, b^\dagger as compared to (6.3a). These just mean that we have 26 copies of isomorphic Moyal algebras⁴. These identifications requires the conditions

$$\mu^{11}(\kappa) = \frac{\theta_\kappa^2 - 4}{\theta_\kappa^2 + 12}, \quad \mu^{12}(\kappa) + \mu^{21}(\kappa) = \frac{16}{\theta_\kappa^2 + 12}, \quad \mu^{12}(\kappa) - \mu^{21}(\kappa) = \frac{8\theta_\kappa}{\theta_\kappa^2 + 12}; \quad (6.7a)$$

$$\mu_\xi^{11} = \frac{\theta_\xi^2 - 4}{\theta_\xi^2 + 12}, \quad \mu_\xi^{12} + \mu_\xi^{21} = \frac{16}{\theta_\xi^2 + 12}, \quad \mu_\xi^{12} - \mu_\xi^{21} = \frac{8\theta_\xi}{\theta_\xi^2 + 12}. \quad (6.7b)$$

Using the explicit expressions (5.3) we can easily solve these equations and find that

$$\theta_\kappa = 2 \tanh \frac{\pi\kappa}{4} \quad \text{and} \quad \theta_\xi = 2 \coth \frac{\eta(b)}{2}, \quad (6.8)$$

where the function $\eta(b)$ is defined a solution of the equation (3.19) (see also Figure 4). θ_κ and θ_ξ are non-commutative parameters associated to the continuous and discrete Moyal algebras respectively. Notice that for $\kappa = 0$ there is only one nonzero oscillator $e_{\kappa=0}$, and $\theta_{\kappa=0}$ is also zero. Therefore for $\kappa = 0$ the associated Moyal algebra is actually a commutative one.

For continuous Moyal algebra the non-commutativity parameter θ_κ is belongs to the interval $[0, 2)$. For discrete Moyal algebra the non-commutativity parameter θ_ξ lies in the interval $(2, \infty)$. From the equations (6.8) and (3.19) follows that when $b \rightarrow \infty$ (i.e. $\alpha' \rightarrow 0$) the parameter θ_ξ goes to 2. It seems that in the low energy limit of the SFT action we have some non-commuting variables. The analysis of the generating function for eigenvectors $V^{(\xi)}$ shows that it goes to 0 as e^{-e^b} as $b \rightarrow \infty$ (see also Figure 5). Therefore regardless of the fact that θ_ξ is finite for $b \rightarrow \infty$ actually there are no oscillator modes corresponding to this non-commutativity.

7 Discussion

Here I would like to summarize the main results of this paper:

1. We explicitly diagonalize the Neumann matrices defining 3-string vertex in the matter and ghost sectors. The spectrum of these operators is given through the spectrum of the unitary auxiliary operators CU and $C\tilde{U}$ respectively.

⁴ More precisely the oscillators $a_{\alpha,\mu}$ are in the space $T(\mathbb{R}^2) \otimes T(\mathbb{R}^{26})$. The tensor $\Theta^{\alpha\beta}$ acts on the first index (i.e. α), while metric in the string target space $g_{\mu\nu}$ acts on the second one.

- Detailed spectrum of the operator CU from the matter sector can be found in Section 3.3. The spectrum of the matter Neumann matrices have been also found by Bo Feng, Yang-Hui He and Nicolas Moeller [14]. To compare the results let us remind the spectrum of the operator M^{11} (5.3). The spectrum consists of two branches: continuous and discrete. The eigenvalues from continuous spectrum are in the interval $[-1/3, 0)$. Each eigenvalue in the interval $(-1/3, 0)$ is doubly degenerated, while for eigenvalue $-1/3$ corresponds only one eigenvector. The discrete spectrum is doubly degenerated and consist of one point in the interval $(0, 1)$. The position of this point depends on the value of parameter b . Basically, the results presented here and the ones obtained in [14] are the same, except for the point $-1/3$ from the continuous spectrum, for which there is one more eigenvector in [14]. The origin of this difference is in improper normalization [15], [16] of the eigenvectors from continuous spectrum used in [14].
 - The spectrum of operator $C\tilde{U}$ from ghost sector is given in Section 4.
2. Using the diagonal representation of the Neumann matrices in the matter sector we identify the Moyal structures appearing in the matter 3-string vertex (Section 6). There are two types of Moyal algebras appearing here: continuous (found in [6]) and discrete. The deformation parameter of the continuous Moyal algebra is given by continuous function $\theta(\kappa)$ defined in (6.8). The value of the continuous deformation parameter is restricted to the interval $[0, 2)$. The deformation parameter of the discrete Moyal algebra is given by θ_ξ defined in (6.8). The value of the discrete deformation parameter depends on the parameter b and is in the interval $(2, \infty)$. As b goes to infinity (i.e. $\alpha' \rightarrow 0$) θ_ξ goes to 2. The corresponding non-commutative modes goes to zero as e^{-e^b} . Despite of the fact that $\theta_\xi = 2$ for $\alpha' = 0$ the non-noncommutative modes are equal to zero.
 3. Notice that the commuting mode for the full string product is $X(\frac{\pi}{2})$ (see (5.7)), while for the product reduced to the zero momentum sector it is left momentum P_L [6].

Below we list some open questions:

1. Identification of ghost vertex algebra with the known ones [17];
2. In the framework of [6] try to establish the exact correspondence between Witten's string product and Moyal product [17];
3. Rewrite the BRST charge in the diagonal basis;
4. Investigate the sliver in the diagonal basis. Classify all projectors of the open string field algebra.

Acknowledgments

I would like to thank H. Liu, G. Moore and B. Zwiebach for useful discussions, B. Feng for correspondence on the spectrum of operator M^{11} , A. Konechny for discussion of the ghost vertices

and comments on the manuscript. And especially I would like to thank S. Lukyanov for communication at various stages of my work. I would like to acknowledge the hospitality of Lawrence Berkeley National Laboratory, where the final pages of the paper were written. The work was supported in part by RFBR grant 02-01-00695.

Appendix

A Normalization of the coordinate and momentum eigenstates

The momentum and coordinate eigenstates are given by the following formulae

$$|p\rangle = \left[\frac{2\pi}{b\alpha'} \right]^{-\frac{d}{4}} \exp \left[-\frac{b\alpha'}{4}(p, p) + \sqrt{b\alpha'}(a_0^\dagger, p) - \frac{1}{2}(a_0^\dagger, a_0^\dagger) \right] |\Omega_b\rangle, \quad (\text{A.1a})$$

$$|x\rangle = \frac{1}{(2\pi b\alpha')^{d/4}} \exp \left[-\frac{1}{b\alpha'}(x, x) + \frac{2i}{\sqrt{b\alpha'}}(a_0^\dagger, x) + \frac{1}{2}(a_0^\dagger, a_0^\dagger) \right] |\Omega_b\rangle. \quad (\text{A.1b})$$

The scalar product (\cdot, \cdot) and the vacuum state $|\Omega_b\rangle$ are defined by

$$(p, p) \equiv g^{\mu\nu} p_\mu p_\nu, \quad (x, x) \equiv g_{\mu\nu} x^\mu x^\nu, \quad (x, p) \equiv x^\mu p_\mu \quad \text{and} \quad a_n |\Omega_b\rangle = 0 \quad \text{for} \quad n \geq 0, \quad (\text{A.2})$$

where $g_{\mu\nu}$ is constant symmetric non-degenerate $d \times d$ matrix (constant metric) and $g^{\mu\nu}$ is its inverse. The conjugated states to the eigenstates (A.1) are given by

$$\langle p| = \left[\frac{2\pi}{b\alpha'} \right]^{-\frac{d}{4}} \langle \Omega_b| \exp \left[-\frac{b\alpha'}{4}(p, p) - \sqrt{b\alpha'}(a_0, p) - \frac{1}{2}(a_0, a_0) \right] \quad (\text{A.3a})$$

$$\langle x| = \frac{1}{(2\pi b\alpha')^{d/4}} \langle \Omega_b| \exp \left[-\frac{1}{b\alpha'}(x, x) - \frac{2i}{\sqrt{b\alpha'}}(a_0, x) + \frac{1}{2}(a_0, a_0) \right]. \quad (\text{A.3b})$$

Normalization of the momentum and coordinate eigenstates:

$$\langle p|p'\rangle = \sqrt{\det g} \delta^{(d)}(p_\mu + p'_\mu) \quad \text{and} \quad \langle x|x'\rangle = \frac{1}{\sqrt{\det g}} \delta^{(d)}(x^\mu + x'^\mu); \quad (\text{A.4a})$$

normalization of the function $\langle p|x\rangle$:

$$\langle p|x\rangle = \overline{\langle x|p\rangle} = \frac{1}{(2\pi)^{\frac{d}{2}}} e^{ip_\mu x^\mu}; \quad (\text{A.4b})$$

string field:

$$|t\rangle = \int \frac{d^d p}{\sqrt{\det g}} t(p)|p\rangle, \quad t(-p) = \langle p|t\rangle; \quad (\text{A.4c})$$

Fourier transform:

$$t(x) = \frac{1}{(2\pi)^{\frac{d}{2}}} \int \frac{d^d p}{\sqrt{\det g}} e^{-ip_\mu x^\mu} t(p) \quad \text{and} \quad t(p) = \frac{1}{(2\pi)^{\frac{d}{2}}} \int d^d x \sqrt{\det g} e^{ip_\mu x^\mu} t(x). \quad (\text{A.4d})$$

B Conformal Ghosts

B.1 Search for the discrete spectrum

B.1.1 Parametrization of the eigenvalues

First of all, let us express the parameter κ appearing in the (3.10) through ν . So we want to solve the equation

$$(2\nu + 1) \cosh \frac{\pi\kappa}{2} + 2 + \nu = -i\sqrt{3} \sinh \frac{\pi\kappa}{2}.$$

This equation has two solutions:

1. ν -independent solution

$$\cosh \frac{\pi\kappa}{2} = -\frac{1}{2} \quad \text{and} \quad \sinh \frac{\pi\kappa}{2} = i\frac{\sqrt{3}}{2} \quad \Rightarrow \quad \frac{\pi\kappa}{2} = \frac{2\pi i}{3} + 2\pi in. \quad (\text{B.1a})$$

2. ν -dependent solution

$$\cosh \frac{\pi\kappa}{2} = -\frac{1}{2} \frac{\nu^2 + 4\nu + 1}{\nu^2 + \nu + 1} \quad \text{and} \quad \sinh \frac{\pi\kappa}{2} = -i\frac{\sqrt{3}}{2} \frac{\nu^2 - 1}{\nu^2 + \nu + 1}. \quad (\text{B.1b})$$

We will always assume that ν lies on the unit circle (see Figure 1), i.e. $\nu = e^{i\phi}$. It convenient to rewrite the formulae above in the following form

$$\cosh \frac{\pi\kappa}{2} = -\frac{\cos \phi + 2}{2 \cos \phi + 1} \quad \text{and} \quad \sinh \frac{\pi\kappa}{2} = \sqrt{3} \frac{\sin \phi}{2 \cos \phi + 1}. \quad (\text{B.1c})$$

If $\frac{2\pi}{3} \leq \phi \leq \frac{4\pi}{3}$ then we will assume $\frac{\pi\kappa}{2} \in \mathbb{R}$, otherwise if $-\frac{2\pi}{3} \leq \phi \leq \frac{2\pi}{3}$ we will assume $\frac{\pi\kappa}{2} \in \mathbb{R} + \pi i$. Moreover it follows from this parametrization that if κ is a solution for ν then $-\kappa$ is a solution for $\bar{\nu}$.

B.1.2 Computation of the integral

Now we are ready to compute the following integral

$$(\bar{W}, \frac{1}{C'U' - \xi} W) = \int_{-\infty}^{\infty} d\kappa \frac{1}{\mathcal{N}(\kappa)} \frac{[(v^{(\kappa)}, W)]^2}{\nu(\kappa) - \xi} \stackrel{3.13b}{=} 2 \int_{-\infty}^{\infty} \frac{d\kappa}{\kappa} \frac{[2 + \nu(\kappa) + \bar{\nu}(\kappa)]}{\kappa \mathcal{N}(\kappa)} \frac{\nu(\kappa)}{\nu(\kappa) - \xi}. \quad (\text{B.2})$$

Substitution of the expressions (3.10) and (3.11) and some simplification yields

$$(W, \frac{1}{C'U' - \xi} W) = 2 \int_{-\infty}^{\infty} \frac{dx}{x} \frac{\sinh \frac{x}{2}}{\cosh \frac{x}{2}} \frac{1}{(1 + 2 \cosh x)} \frac{2 + \cosh x + i\sqrt{3} \sinh x}{(2\xi + 1) \cosh x + i\sqrt{3} \sinh x + 2 + \xi}, \quad (\text{B.3})$$

where $x = \frac{\pi\kappa}{2}$. The analysis performed in (B.1) yields to the following parametrization of the ξ :

$$\cosh(\eta + i\pi) = -\frac{1}{2} \frac{\xi^2 + 4\xi + 1}{\xi^2 + \xi + 1} \quad \text{and} \quad \sinh(\eta + i\pi) = -i\frac{\sqrt{3}}{2} \frac{\xi^2 - 1}{\xi^2 + \xi + 1}. \quad (\text{B.4})$$

The integral (B.3) has the following poles (see Figure 3):




Figure 3: Contour of integration for expression (B.3).

- $x = \pi i + 2\pi i n, n \geq 0$;
- $x = \eta + i\pi + 2\pi i n, n \geq 0$;
- $x = \frac{4\pi i}{3} + 2\pi i n, n \geq 0$.

One sees that we have two cases $\eta = 0$ and $\eta \neq 0$.

Let us first consider the case $\eta \neq 0$. The computation of the residues in the upper half plane leads to the following result

$$\begin{aligned}
 (\overline{W}, \frac{1}{C'U' - \xi} W) &= 2 \sum_{n=0}^{\infty} \left[\frac{2}{\xi - 1} \frac{1}{n + \frac{1}{2}} + \frac{1}{n + \frac{2}{3}} - \frac{\xi + 1}{\xi - 1} \frac{1}{n + \frac{1}{2} + \frac{\eta}{2\pi i}} \right] \\
 &= \frac{\xi + 1}{\xi - 1} 2F(\eta) + 2[\psi(\frac{1}{2}) - \psi(\frac{2}{3})], \quad (\text{B.5})
 \end{aligned}$$

where $\psi(x)$ is the logarithmic derivative of the Euler Γ -function

$$\begin{aligned}
 \psi(x) &= -\gamma_E - \sum_{n=0}^{\infty} \left(\frac{1}{n+x} - \frac{1}{n+1} \right), \\
 \psi(\frac{1}{2}) &= -\gamma_E - \log 4 \quad \text{and} \quad \psi(\frac{2}{3}) = -\gamma_E + \frac{\pi}{2\sqrt{3}} - \log 3\sqrt{3}
 \end{aligned}$$

and

$$F(\eta) = \psi(\frac{1}{2} + \frac{\eta}{2\pi i}) - \psi(\frac{1}{2}). \quad (\text{B.6})$$

If $\eta = 0$ we need only to know that the integral (B.3) is finite. It follows from equation (3.18b) that $\eta = 0$ and $\xi = 1$ is its solution only for $b = 0$.




Figure 4: On the left pane we present the graph of function $\Re F(\eta)$. As η goes to ∞ this function grows logarithmically. On the right pane we present the graph of $\arg \bar{\xi}(b)$. As b grows this function goes to its maximal value $\frac{2\pi}{3}$ exponentially fast.

B.1.3 Solving of the equation

Let us simplify expression (B.5). To this end we substitute $\psi(\frac{1}{2})$ and $\psi(\frac{2}{3})$ and notice that⁵

$$\Im F(\eta) = -\frac{\pi}{2} \tanh \frac{\eta}{2} \equiv \frac{\pi}{2i\sqrt{3}} \frac{\xi - 1}{\xi + 1}. \quad (\text{B.7})$$

Therefore (B.5) takes a form

$$(\bar{W}, \frac{1}{C'U' - \xi} W) = V_{00} + \frac{\xi + 1}{\xi - 1} 2\Re F(\eta). \quad (\text{B.8})$$

Substitution of this expression into equation (3.18b) yields the equation

$$\Re F(\eta) = \frac{b}{4}. \quad (\text{B.9})$$

$\Re F(\eta)$ is an even function with $F(0) = 0$ (see Figure 4), therefore for each $b > 0$ this equation has two solutions $\pm\eta(b)$. On Figure 4 we presented the $\arg \bar{\xi}(b)$ for this solution.

⁵This expression follows from the following formula for the Γ -function

$$\Gamma(1 - z)\Gamma(z) = \frac{\pi}{\sin \pi z}.$$

B.2 Generating function for discrete spectrum

In this section we will construct the generating function for the vector defined by (3.18a). To this end we rewrite the expression (3.18a) for the components of the vector $V^{(\xi)}$ in the following form

$$V_m^{(\xi)} = \frac{V_0^{(\xi)}}{\sqrt{b}}(\xi - 1) \left(\frac{1}{C'U' - \xi} \right)_{mn} W_n \stackrel{3.12}{=} \frac{V_0^{(\xi)}}{\sqrt{b}}(\xi - 1) \int_{-\infty}^{\infty} \frac{d\kappa}{\mathcal{N}(\kappa)} v_m^{(\kappa)} \left(v^{(\kappa)}, \frac{1}{C'U' - \xi} W \right) \\ \stackrel{3.13b}{=} -\frac{V_0^{(\xi)}}{\sqrt{b}}(\xi - 1) \int_{-\infty}^{\infty} \frac{d\kappa}{\kappa \mathcal{N}(\kappa)} v_m^{(\kappa)} \frac{\sqrt{2}[1 + \nu(\kappa)]}{\nu(\kappa) - \xi}.$$

The generating function for the vector $V_m^{(\xi)}$ is defined as

$$F^{(\xi)}(z) = \sum_{m=1}^{\infty} V_m^{(\xi)} \frac{z^m}{\sqrt{m}} = -\sqrt{2} \frac{V_0^{(\xi)}}{\sqrt{b}}(\xi - 1) \int_{-\infty}^{\infty} \frac{d\kappa}{\kappa \mathcal{N}(\kappa)} \frac{f^{(\kappa)}(z)[1 + \nu(\kappa)]}{\nu(\kappa) - \xi}. \quad (\text{B.10})$$

To get the explicit expression for this generating function we need to compute the integral

$$\int_{-\infty}^{\infty} \frac{d\kappa}{\kappa \mathcal{N}(\kappa)} \frac{f^{(\kappa)}(z)[1 + \nu(\kappa)]}{\nu(\kappa) - \xi} \stackrel{3.10}{=} \int_{-\infty}^{\infty} \frac{dx}{x} \frac{1 - e^{-x \frac{2}{\pi} \arctan z}}{2 \sinh x} \frac{1 - \cosh x + i\sqrt{3} \sinh x}{(2\xi + 1) \cosh x + i\sqrt{3} \sinh x + 2 + \xi}$$

We want to compute this integral by summing residues in the UHP. To this end we add $+ix\delta$ ($\delta > 0$) to the exponent. This integral has the following poles

- $x = \pi i + 2\pi i n, n \geq 0;$
- $x = \eta + \pi i + 2\pi i n, n \geq 0.$

Summing over these residues yields

$$= -\frac{1}{\xi - 1} \sum_{n=0}^{\infty} \left[-\frac{1}{n + \frac{1}{2}} + \frac{e^{-4i(n + \frac{1}{2})(\arctan z - i\delta)}}{n + \frac{1}{2}} + \frac{1}{n + \frac{1}{2} + \frac{\eta}{2\pi i}} - \frac{e^{-4i(n + \frac{1}{2} + \frac{\eta}{2\pi i})(\arctan z - i\delta)}}{n + \frac{1}{2} + \frac{\eta}{2\pi i}} \right] \\ = \frac{1}{\xi - 1} \left[\psi\left(\frac{1}{2} + \frac{\eta}{2\pi i}\right) - \psi\left(\frac{1}{2}\right) + e^{-2i(1 + \frac{\eta}{\pi i}) \arctan z} \text{LerchPhi}\left(e^{-4i \arctan z}, 1, \frac{1}{2} + \frac{\eta}{2\pi i}\right) \right. \\ \left. - e^{-2i \arctan z} \text{LerchPhi}\left(e^{-4i \arctan z}, 1, \frac{1}{2}\right) \right], \quad (\text{B.11})$$

where

$$\text{LerchPhi}(w, 1, x) = \sum_{n=0}^{\infty} \frac{w^n}{n + x} = \frac{1}{x} {}_2F_1(1, x; x + 1; w).$$

Using the fact that

$${}_2F_1\left(1, \frac{1}{2}; \frac{3}{2}; w\right) = \frac{1}{\sqrt{w}} \operatorname{arctanh} \sqrt{w}, \quad \operatorname{arctanh} e^{-4i \arctan z} = -\frac{1}{2} \log iz$$

and the expressions (B.7), (B.9) we can write the following formula for the generating function $F^{(\xi)}(z)$

$$F^{(\xi)}(z) = -\sqrt{\frac{2}{b}} V_0^{(\xi)} \left[\frac{b}{4} + \frac{\pi}{2\sqrt{3}} \frac{\xi - 1}{\xi + 1} + \log iz + e^{-2i(1+\frac{\eta}{\pi i}) \arctan z} \text{LerchPhi}(e^{-4i \arctan z}, 1, \frac{1}{2} + \frac{\eta}{2\pi i}) \right], \quad (\text{B.12})$$

where $F(\eta)$ is defined by (B.6) and η is related to ξ by formula (B.4).

Having these formulae one can easily calculate how complex conjugation and operator C change the eigenvectors:

$$F^{(\xi)}(-z) \equiv F^{(\bar{\xi})}(z) \quad \text{and} \quad \overline{F^{(\xi)}(z)} \equiv F^{(\xi)}(z) \quad \text{iff} \quad V_0^{(\xi)} \in \mathbb{R}. \quad (\text{B.13})$$

B.3 Normalization of the vectors from the discrete spectrum

In this subsection we will compute the following scalar product

$$((V^{(\xi)}, V^{(\xi)})) := |V_0^{(\xi)}|^2 + (V^{(\xi)}, V^{(\xi)}). \quad (\text{B.14})$$

The expression (B.14) yields

$$\begin{aligned} (V^{(\xi)}, V^{(\xi)}) &\stackrel{3.18a}{=} \frac{|V_0^{(\xi)}|^2}{b} |\xi - 1|^2 \left(W, \frac{1}{U' C' - \bar{\xi} C' U' - \xi} W \right) \\ &\stackrel{3.12, 3.13b}{=} \frac{|V_0^{(\xi)}|^2}{b} |\xi - 1|^2 2 \int_{-\infty}^{\infty} \frac{d\kappa}{\kappa} \frac{2 + \nu + \bar{\nu}}{\kappa \mathcal{N}(\kappa)} \frac{1}{(\nu - \xi)(\bar{\nu} - \bar{\xi})} \\ &= \frac{|V_0^{(\xi)}|^2}{b} |\xi - 1|^2 (-\xi) \frac{\partial}{\partial \xi} \left[2 \int_{-\infty}^{\infty} \frac{d\kappa}{\kappa} \frac{2 + \nu + \bar{\nu}}{\kappa \mathcal{N}(\kappa)} \frac{\nu}{\nu - \xi} \right] \\ &\stackrel{B.2, B.8}{=} \frac{|V_0^{(\xi)}|^2}{b} |\xi - 1|^2 \left[-\frac{b}{|\xi - 1|^2} + \frac{2\sqrt{3}}{i} \frac{\xi + 1}{\xi - 1} \frac{1}{\xi + \bar{\xi} + 1} \frac{\partial}{\partial \eta} \Re F(\eta) \right]. \quad (\text{B.15}) \end{aligned}$$

Substitution of this expression into (B.14) and simplification by use of (B.9) yields

$$((V^{(\xi)}, V^{(\xi)})) = |V_0^{(\xi)}|^2 \sinh \eta \frac{\partial}{\partial \eta} \left[\log \Re F(\eta) \right]. \quad (\text{B.16})$$

The function on the rhs is positive and more or equal to 2. So to have the norm equal to one, we have to choose $V_0^{(\xi)}$ equal to

$$V_0^{(\xi)} = \left(\sinh \eta \frac{\partial}{\partial \eta} \left[\log \Re F(\eta) \right] \right)^{-1/2}. \quad (\text{B.17})$$

The graphic of this function is presented on the Figure 5. Notice that as $\eta \rightarrow \infty$ $V_0^{(\xi)}$ goes to zero exponentially fast.




Figure 5: Plot of the zero component $V_0^{(\xi)}$ of the vectors from continuous spectra. If we choose zero component in this way the vectors will be normalized to 1.

B.4 Generating function for the continuous spectrum

B.4.1 Calculation of $A^{(\kappa)}$

First of all we need to obtain the expression for the $A^{(\kappa)}$ defined by equation (3.17). To this end we have to calculate the integral

$$\begin{aligned} \left(\overline{W}, \mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W \right) &= 2 \int_{-\infty}^{\infty} \frac{d\kappa'}{\kappa'} \frac{2 + \nu + \bar{\nu}}{\kappa \mathcal{N}(\kappa')} \mathcal{P} \frac{\nu(\kappa')}{\nu(\kappa') - \nu(\kappa)} \\ &= 2 \int \frac{dx \sinh \frac{x}{2}}{x \cosh \frac{x}{2}} \frac{1}{1 + 2 \cosh x} \mathcal{P} \frac{2 + \cosh x + i\sqrt{3} \sinh x}{(2\nu + 1) \cosh x + i\sqrt{3} \sinh x + 2 + \nu}. \end{aligned} \quad (\text{B.18})$$

This integral has the following poles:

- $x = \pi i + 2\pi i n, n \geq 0$;
- $x = \frac{\pi \kappa}{2} + 2\pi i(n + 1), n \geq 0$;
- $x = \frac{\pi \kappa}{2}$, one half of the residue;
- $x = \frac{4\pi i}{3} + 2\pi i n, n \geq 0$.

Calculation of the residues yields

$$\begin{aligned} &= 2 \sum_{n=0}^{\infty} \left[\frac{2}{\nu - 1} \frac{1}{n + \frac{1}{2}} - \frac{\nu + 1}{\nu - 1} \frac{1}{n + 1 + \frac{\kappa}{4i}} + \frac{1}{n + \frac{2}{3}} \right] - \frac{\nu + 1}{\nu - 1} \frac{4i}{\kappa} \\ &= -\frac{\pi}{\sqrt{3}} + V_{00} + 2 \frac{\nu + 1}{\nu - 1} \left[\psi\left(1 + \frac{\kappa}{4i}\right) - \psi\left(\frac{1}{2}\right) - \frac{2i}{\kappa} \right] \\ &= V_{00} + \frac{\nu + 1}{\nu - 1} 2\Re F_c(\kappa), \end{aligned} \quad (\text{B.19})$$

where

$$F_c(\kappa) = \psi\left(1 + \frac{\kappa}{4i}\right) - \psi\left(\frac{1}{2}\right) \quad (\text{B.20})$$

and to obtain the last line in the (B.19) we have used the fact that

$$\Im F_c(\kappa) = \frac{2}{\kappa} - \frac{\pi \sinh \frac{\pi\kappa}{2}}{2 \cosh \frac{\pi\kappa}{2}} \equiv \frac{2}{\kappa} - \frac{\pi i}{2\sqrt{3}} \frac{\nu - 1}{\nu + 1}.$$

Substitution of (B.19) into (3.17) yields the following expression for $A^{(\kappa)}$:

$$A^{(\kappa)} = V_0^{(\kappa)} \sqrt{\frac{2}{b}} \kappa \left(\Re F_c(\kappa) - \frac{b}{4} \right). \quad (\text{B.21})$$

B.4.2 Generating function

The generating function for the vector $V_m^{(\kappa)}$ is defined in the following way

$$\begin{aligned} F^{(\kappa)}(z) &= \sum_{m=1}^{\infty} V_m^{(\kappa)} \frac{z^m}{\sqrt{m}} \\ &\stackrel{3.17a}{=} A^{(\kappa)} f^{(\kappa)}(z) - [1 - \nu(\kappa)] \frac{V_0^{(\kappa)}}{\sqrt{b}} \int \frac{d\kappa'}{\mathcal{N}(\kappa')} f^{(\kappa')}(z) \left(v^{(\kappa')}, \mathcal{P} \frac{1}{CU - \nu(\kappa)} W \right). \end{aligned} \quad (\text{B.22})$$

Let us start from the computing the integral in the rhs:

$$\begin{aligned} \int \frac{d\kappa'}{\mathcal{N}(\kappa')} f^{(\kappa')}(z) \mathcal{P} \frac{(v^{(\kappa')}, W)}{\nu(\kappa') - \nu(\kappa)} &= -\sqrt{2} \int \frac{d\kappa'}{\kappa' \mathcal{N}(\kappa')} \frac{1}{\kappa'} (1 - e^{-\kappa \arctan z}) \mathcal{P} \frac{1 + \nu(\kappa')}{\nu(\kappa') - \nu(\kappa)} \\ &= -\sqrt{2} \int_{-\infty}^{\infty} \frac{dx}{x} \frac{1 - e^{-x \frac{2}{\pi} \arctan z}}{2 \sinh x} \mathcal{P} \frac{1 - \cosh x + i\sqrt{3} \sinh x}{(2\nu + 1) \cosh x + i\sqrt{3} \sinh x + 2 + \nu}. \end{aligned} \quad (\text{B.23})$$

This integral has the following poles

- $x = \pi i + 2\pi n$, $n \geq 0$;
- $x = \frac{\pi\kappa}{2} + 2\pi(n + 1)$, $n \geq 0$;
- $x = \frac{\pi\kappa}{2}$, half of the pole.

Calculation of the residues yields

$$\begin{aligned}
&= -\frac{\sqrt{2}}{1-\nu} \sum_{n=0}^{\infty} \left[-\frac{1 - e^{-(n+\frac{1}{2})4i \arctan z}}{n + \frac{1}{2}} + \frac{1 - e^{-(n+1+\frac{\kappa}{4i})4i \arctan z}}{n + 1 + \frac{\kappa}{4i}} \right] - \frac{\sqrt{2}}{1-\nu} \frac{2i}{\kappa} (1 - e^{-\kappa \arctan z}) \\
&= -\frac{\sqrt{2}}{1-\nu} \left[\psi\left(\frac{1}{2}\right) - \psi\left(1 + \frac{\kappa}{4i}\right) + \frac{2i}{\kappa} (1 - e^{-\kappa \arctan z}) + \text{LerchPhi}\left(e^{-4i \arctan z}, 1, \frac{1}{2}\right) e^{-2i \arctan z} \right. \\
&\quad \left. - \text{LerchPhi}\left(e^{-4i \arctan z}, 1, 1 + \frac{\kappa}{4i}\right) e^{-4i \arctan z} e^{-\kappa \arctan z} \right] \\
&= \frac{\sqrt{2}}{1-\nu} \left[\Re F_c(\kappa) + \left(\frac{\pi}{2\sqrt{3}} \frac{\nu-1}{\nu+1} + \frac{2i}{\kappa} \right) + \log iz - 2i f^{(\kappa)}(z) \right. \\
&\quad \left. + \text{LerchPhi}\left(e^{-4i \arctan z}, 1, 1 + \frac{\kappa}{4i}\right) e^{-4i \arctan z} e^{-\kappa \arctan z} \right]. \quad (\text{B.24})
\end{aligned}$$

Substitution of this expression into (B.22) yields

$$\begin{aligned}
F^{(\kappa)}(z) &= V_0^{(\kappa)} \left[\frac{2}{b} \right]^{\frac{1}{2}} \left[-\frac{b}{4} - \left(\Re F_c(\kappa) - \frac{b}{4} \right) e^{-\kappa \arctan z} - \left(\frac{\pi}{2\sqrt{3}} \frac{\nu-1}{\nu+1} + \frac{2i}{\kappa} \right) - \log iz + 2i f^{(\kappa)}(z) \right. \\
&\quad \left. - \text{LerchPhi}\left(e^{-4i \arctan z}, 1, 1 + \frac{\kappa}{4i}\right) e^{-4i \arctan z} e^{-\kappa \arctan z} \right]. \quad (\text{B.25})
\end{aligned}$$

Now let us consider the generating function for the $\kappa = 0$. Using the fact that

$$\text{LerchPhi}(w, 1, 1) = {}_2F_1(1, 1; 2; w) = -\frac{\log(1-w)}{w}$$

one obtains the following expression

$$F^{(0)}(z) = -V_0^{(0)} \left[\frac{2}{b} \right]^{\frac{1}{2}} \log(1+z^2). \quad (\text{B.26})$$

B.5 Normalization of the vectors from continuous spectrum

The aim of this subsection is to compute the following scalar product

$$((V^{(\kappa)}, V^{(\kappa')})) = \overline{V}_0^{(\kappa)} V_0^{(\kappa')} + (V^{(\kappa)}, V^{(\kappa')}). \quad (\text{B.27})$$

Substitution of (3.17a) yields

$$\begin{aligned}
((V^{(\kappa)}, V^{(\kappa')})) &= \overline{V}_0^{(\kappa)} V_0^{(\kappa')} + \overline{A^{(\kappa)}} A^{(\kappa')} (v^{(\kappa)}, v^{(\kappa')}) - [1 - \overline{\nu}(\kappa)] \frac{\overline{V}_0^{(\kappa)} A^{(\kappa')}}{\sqrt{b}} \left(\mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W, v^{(\kappa')} \right) \\
&\quad - [1 - \nu(\kappa')] \frac{V_0^{(\kappa')} \overline{A^{(\kappa')}}}{\sqrt{b}} \left(v^{(\kappa)}, \mathcal{P} \frac{1}{C'U' - \nu(\kappa')} W \right) \\
&\quad + [1 - \overline{\nu}(\kappa)] [1 - \nu(\kappa')] \frac{V_0^{(\kappa')} \overline{V}_0^{(\kappa)}}{b} \left(\mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W, \mathcal{P} \frac{1}{C'U' - \nu(\kappa')} W \right).
\end{aligned}$$

Substitution of the expressions (3.11), (3.13b) and (B.21) yields

$$\begin{aligned}
&= |V_0^{(\kappa)}|^2 \mathcal{N}(\kappa) \frac{2}{b} \kappa^2 \left[\Re F_c(\kappa) - \frac{b}{4} \right]^2 \delta(\kappa - \kappa') + \frac{2}{b} \frac{\bar{V}_0^{(\kappa)} V_0^{(\kappa')}}{\nu(\kappa) - \nu(\kappa')} \left[\Re F_c(\kappa') (\nu(\kappa) - 1)(\nu(\kappa') + 1) - (\kappa \leftrightarrow \kappa') \right] \\
&\quad + [1 - \bar{\nu}(\kappa)][1 - \nu(\kappa')] \frac{\bar{V}_0^{(\kappa)} V_0^{(\kappa')}}{b} \left(\mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W, \mathcal{P} \frac{1}{C'U' - \nu(\kappa')} W \right). \quad (\text{B.28})
\end{aligned}$$

Let us now compute the expression

$$\begin{aligned}
&\left(\mathcal{P} \frac{1}{C'U' - \nu(\kappa)} W, \mathcal{P} \frac{1}{C'U' - \nu(\kappa')} W \right) \\
&= -2\nu(\kappa) \int_{-\infty}^{\infty} \frac{d\kappa''}{\kappa''} \frac{[1 + \nu(\kappa'')]^2}{\kappa'' \mathcal{N}(\kappa'')} \mathcal{P} \frac{1}{\nu(\kappa'') - \nu(\kappa')} \mathcal{P} \frac{1}{\nu(\kappa'') - \nu(\kappa)}. \quad (\text{B.29})
\end{aligned}$$

We have to compute this integral very carefully, because it has a singularity then $\kappa' = \kappa$. To extract it we will use the following identity

$$\mathcal{P} \frac{1}{x'' - x} = \frac{1}{x'' - x + i0} + \pi i \delta(x'' - x).$$

In our case this relation takes a form

$$\mathcal{P} \frac{1}{\nu(\kappa'') - \nu(\kappa)} = \frac{1}{\nu(\kappa'') - \nu(\kappa) + i0} + \frac{\pi i}{\frac{d\nu}{d\kappa}} \delta(\kappa'' - \kappa).$$

In principal, one has to understand all these relations as a specification of the integration contour in (B.29). Substitution of this relation into (B.29) yields

$$\begin{aligned}
&= -2\nu(\kappa) \int_{-\infty}^{\infty} \frac{d\kappa''}{\kappa''} \frac{[1 + \nu(\kappa'')]^2}{\kappa'' \mathcal{N}(\kappa'')} \left[\frac{1}{\nu(\kappa'') - \nu(\kappa') + i0} \frac{1}{\nu(\kappa'') - \nu(\kappa) + i0} \right. \\
&\quad + \frac{\pi i}{\frac{d\nu}{d\kappa}} \delta(\kappa'' - \kappa) \frac{1}{\nu(\kappa'') - \nu(\kappa') + i0} + \frac{\pi i}{\frac{d\nu}{d\kappa'}} \delta(\kappa'' - \kappa') \frac{1}{\nu(\kappa'') - \nu(\kappa) + i0} \\
&\quad \left. - \frac{\pi^2}{\frac{d\nu}{d\kappa} \frac{d\nu}{d\kappa'}} \delta(\kappa'' - \kappa) \delta(\kappa'' - \kappa') \right]
\end{aligned}$$

Using the relation

$$\frac{2\kappa}{\pi} \mathcal{N}(\kappa) \frac{d\nu(\kappa)}{d\kappa} = [1 - \nu(\kappa)]^2$$

we can simplify this expression

$$\begin{aligned}
&= -2\nu(\kappa) \int_{-\infty}^{\infty} \frac{d\kappa''}{\kappa''} \frac{[1 + \nu(\kappa'')]^2}{\kappa'' \mathcal{N}(\kappa'')} \left[\frac{1}{\nu(\kappa'') - \nu(\kappa') + i0} \frac{1}{\nu(\kappa'') - \nu(\kappa) + i0} \right] \\
&\quad + \mathcal{P} \frac{\nu(\kappa)}{\nu(\kappa) - \nu(\kappa')} \left[\frac{4i}{\kappa'} \frac{1 + \nu(\kappa')}{1 - \nu(\kappa')} - \frac{4i}{\kappa} \frac{1 + \nu(\kappa)}{1 - \nu(\kappa)} \right] + \frac{8\mathcal{N}(\kappa)}{[1 - \nu(\kappa)][1 - \bar{\nu}(\kappa)]} \delta(\kappa - \kappa'). \quad (\text{B.30})
\end{aligned}$$

The integral in the first line can be rewritten as

$$2 \int_{-\infty}^{\infty} \frac{dx'' \sinh \frac{x''}{2}}{x'' \cosh \frac{x''}{2}} \frac{1 + 2 \cosh x''}{[(2\nu' + 1) \cosh x'' + i\sqrt{3} \sinh x'' + 2 + \nu' + i0]} \times \frac{1}{(2\nu + 1) \cosh x'' + i\sqrt{3} \sinh x'' + 2 + \nu + i0}.$$

This integral has the following poles

- $x'' = \pi i + 2\pi i n$, for $n \geq 0$;
- $x'' = \frac{\pi \kappa'}{2} + 2\pi i n$, for $n \geq 1$;
- $x'' = -\frac{\pi \kappa'}{2} + 2\pi i n$, for $n \geq 1$.

Calculation of the residues yields the following expression for (B.30)

$$\begin{aligned} &= 2 \sum_{n=0}^{\infty} \left[\frac{1}{n + \frac{1}{2}} \frac{-2}{(1 - \nu')(1 - \nu)} + \mathcal{P} \frac{\nu}{\nu - \nu'} \left(\frac{1 + \nu'}{1 - \nu'} \frac{1}{n + 1 + \frac{\kappa'}{4i}} - \frac{1 + \nu}{1 - \nu} \frac{1}{n + 1 + \frac{\kappa}{4i}} \right) \right] \\ &\quad + \mathcal{P} \frac{\nu}{\nu - \nu'} \left(\frac{1 + \nu'}{1 - \nu'} \frac{4i}{\kappa'} - \frac{1 + \nu}{1 - \nu} \frac{4i}{\kappa} \right) + \frac{8\mathcal{N}(\kappa)}{[1 - \nu(\kappa)][1 - \bar{\nu}(\kappa)]} \delta(\kappa - \kappa'). \\ &= \mathcal{P} \frac{2\nu}{\nu - \nu'} \left[\frac{1 + \nu}{1 - \nu} \Re F_c(\kappa) - \frac{1 + \nu'}{1 - \nu'} \Re F_c(\kappa') \right] + \frac{8\mathcal{N}(\kappa)}{(1 - \nu)(1 - \bar{\nu})} \delta(\kappa - \kappa'). \quad (\text{B.31}) \end{aligned}$$

Substitution of this expression into (B.28) yields

$$((V^{(\kappa)}, V^{(\kappa')})) = \frac{2}{b} |V_0^{(\kappa)}|^2 \mathcal{N}(\kappa) \left[4 + \kappa^2 \left(\Re F_c(\kappa) - \frac{b}{4} \right)^2 \right] \delta(\kappa - \kappa'). \quad (\text{B.32})$$

To obtain standard normalization of the continuous spectra one has to put

$$V_0^{(\kappa)} = \left[\frac{b}{2} \right]^{\frac{1}{2}} \left[\mathcal{N}(\kappa) \right]^{-\frac{1}{2}} \left[4 + \kappa^2 \left(\Re F_c(\kappa) - \frac{b}{4} \right)^2 \right]^{-\frac{1}{2}}. \quad (\text{B.33})$$

References

- [1] E. Witten, “Noncommutative Geometry And String Field Theory,” Nucl. Phys. B **268**, 253 (1986).
- [2] A. Sen, *Stable non BPS bound states of bps d-branes*, JHEP **08**, 010 (1998), hep-th/9805019
A. Sen, *SO(32) spinors of type i and other solitons on brane - anti-brane pair*, JHEP **09**, 023 (1998), hep-th/9808141
- [3] K. Ohmori, *A Review on Tachyon Condensation in Open String Field Theories*, hep-th/0102085.
- [4] P. De Smet, *Tachyon Condensation: Calculations in String Field Theory*, hep-th/0109182.
- [5] I.Ya. Aref’eva, D.M. Belov, A.A. Giryvets, A.S. Koshelev, P.B. Medvedev, *Noncommutative Field Theories and (Super)String Field Theories*, hep-th/0111208.
- [6] M. Douglas, H. Liu, G. Moore and B. Zwiebach, *Open String Star as a Continuous Moyal Product*, hep-th/0202087
- [7] I. Bars, “Map of Witten’s $*$ to Moyal’s $*$,” Phys. Lett. B **517**, 436 (2001), hep-th/0106157.
- [8] D. Gross, A. Jevicki, *Operator Formulation of Interacting String Field Theory (I)*, Nucl.Phys. **B283** (1987) 1 – 49
- [9] D. Gross, A. Jevicki, *Operator Formulation of Interacting String Field Theory (II)*, Nucl.Phys. **B287** (1987) 225 – 250
- [10] L. Rastelli, A. Sen, B. Zwiebach, *Star Algebra Spectroscopy*, hep-th/0111281
- [11] L. Rastelli, A. Sen, B. Zwiebach, *Classical Solutions in String Field Theory Around the Tachyon Vacuum*, hep-th/0102112
- [12] K. Okuyama, *Ghost Kinetic Operator of Vacuum String Field Theory*, hep-th/0201015
- [13] K. Okuyama, *Ratio of Tensions from Vacuum String Field Theory*, hep-th/0201136
- [14] B. Feng, Y.-H. He, N. Moeller, *The Spectrum of the Neumann Matrix with Zero Modes*, hep-th/0202176
- [15] B. Feng, private communication
- [16] B. Feng, Y. He, N. Moeller, *Zeeman Spectroscopy of the Star Algebra*, hep-th/0203175
- [17] D. Belov and A. Konechny, work in progress