arXiv:math/9411205v1 [math.LO] 1 Nov 1994

Souslin's Hypothesis and Convergence in Category

by Arnold W. Miller[1]

Suppose that $S \subseteq P(X)$ is a σ-field of subsets of X and $I \subseteq S$ is a σ-ideal. If I has the countable chain condition (ccc), i.e., every family of disjoint sets in $S \setminus I$ is countable, then S/I is a complete boolean algebra. A boolean algebra is atomic iff there is an atom beneath every nonzero element.

A function $f : X \to \mathbb{R}$ is S-measurable iff $f^{-1}(U) \in S$ for every open set U. A sequence of S-measurable functions $f_n : X \to \mathbb{R}$ converges I-a.e. to a function f iff there exists $A \in I$ such that $f_n(x) \to f(x)$ for all $x \in (X \setminus A)$. If (X, S, μ) is a finite measure space, then a sequence of measurable functions $f_n : X \to \mathbb{R}$ converges in measure to a function f iff for any $\epsilon > 0$ there exists N such that for any $n > N$:

$$\mu(\{x \in X : |f_n(x) - f(x)| > \epsilon\}) < \epsilon.$$

In this case if I is the ideal of measure zero sets, then f_n converges to f in measure iff every subsequence $\{f_n : n \in X\}$ (where $X \subseteq \mathbb{N}$) has a subsequence $Y \subseteq X$ such that $\{f_n : n \in Y\}$ converges I-a.e. This allows us to define convergence in measure without mentioning the measure, only the ideal I. So in the abstract setting define f_n converges to f with respect to I iff every subsequence $\{f_n : n \in X\}$ has a subsequence $Y \subseteq X$ such that $\{f_n : n \in Y\}$ converges I-a.e. (where X and Y range over infinite sets of natural numbers.) For more background on this subject in case I is the ideal of meager sets, see Poreda, Wagner-Bojakoska, and Wilczyński [PWW] and Ciesielski, Larson, and Ostaszewski [CLO].

Marczewski [M] showed that if (X, S, μ) is an atomic measure and I the μ-null sets, then 'I-a.e. convergence' is the same as 'convergence with respect to I'.

[1] I want to thank Krzysztof Ciesielski for many helpful conversations.

The results presented in this paper were obtained during the Joint US–Polish Workshop in Real Analysis, Łódź, Poland, July 1994. The Workshop was partially supported by the NSF grant INT–9401673.

AMS Subject Classification. Primary: 28A20; Secondary: 03E65, 54E52.

Gribanov [G] proved the converse, if (X, S, μ) is a finite measure space and I the μ-null sets, then if 'I-a.e. convergence' is the same as 'convergence with respect to I' then μ is an atomic measure.

Souslin's Hypothesis (SH) is the statement that there are no Souslin lines. It is known to be independent (see Solovay and Tennenbaum [ST]). It was the inspiration for Martin's Axiom.

Theorem 1 *(Wagner and Wilczyński [WW]) Assume SH. Then for any σ-field S and ccc σ-ideal $I \subseteq S$ the following are equivalent:*

- *'I-a.e. convergence' is the same as 'convergence with respect to I' for S-measurable sequences sequences of real-valued functions, and*
- *the complete boolean algebra S/I is atomic.*

At the real analysis meeting in Łódź Poland in July 94, Wilczyński asked whether or not SH is needed for the Theorem above. We show here that the conclusion of Theorem 1 implies Souslin's Hypothesis.

Theorem 2 *Suppose SH is false (so there exists a Souslin tree). Then there exists a regular topological space X such that*

1. *X has no isolated points,*
2. *X is ccc (every family of disjoint open sets is countable),*
3. *every nonempty open subset of X is nonmeager, and*
4. *if I is the σ-ideal of meager subsets of X, then 'I-a.e. convergence' is the same as 'convergence with respect to I' for any sequence of Baire measurable real-valued functions.*

Hence if S is σ-ideal of sets with the property of Baire and I the σ-ideal of meager sets, then S/I is ccc and nonatomic, but the two types of convergence are the same.

Proof: Define $(T, <)$ is an ω_1-tree iff it is a partial order and for each $s \in T$ the set $\{t \in T : t < s\}$ is well-ordered by $<$ with some countable order type, $\alpha < \omega_1$. We let

$$T_\alpha = \{s \in T : \{t \in T : t < s\} \text{ has order type } \alpha\}.$$

Also

$$T_{<\alpha} = \bigcup\{T_\beta : \beta < \alpha\}.$$

Define $C \subseteq T$ is a chain iff for every $s, t \in C$ either $s \leq t$ or $t \leq s$.

Define $A \subseteq T$ is an antichain iff for any $s, t \in A$ if $s \leq t$, then $s = t$, i.e. distinct elements are $\leq$-incomparable.

Define T is a Souslin tree iff T is an ω_1 tree in which every chain and antichain is countable. (Note that since T_α is an antichain it must be countable.)

SH is equivalent to saying there is no Souslin tree. Every Souslin tree contains a normal Souslin tree, i.e., a Souslin tree T such that for every $\alpha < \beta < \omega_1$ and $s \in T_\alpha$ there exists a $t \in T_\beta$ with $s < t$. (Just throw out nodes of T which do not have extensions arbitrarily high in the tree.) For more on Souslin trees see Todorčevič [T].

Now we are ready to define our space X. Let the elements of X be maximal chains of T. For each $s \in T$ let

$$C_s = \{b \in X : s \in b\}$$

and let

$$\{C_s : s \in T\}$$

be an open basis for the topology on X. Note that $C_s \cap C_t$ is either empty or equal to either C_s or C_t depending on whether s and t are incomparable, or $t \leq s$ or $s \leq t$, respectively. Each C_s is clopen since its complement is the union of C_t for t which are incomparable to s. X has no isolated points, since given any $s \in T$ there must be incomparable extensions of s (because T is normal) and therefore at least two maximal chains containing s, so C_s is not a singleton. Clearly X has the countable chain condition.

Lemma 3 *Open subsets of X are nonmeager. In fact, the intersection of countably many open dense sets contains an open dense set.*

Proof: Suppose $(U_n : n \in \omega)$ is a sequence of an open dense subsets of X. Let $A_n \subseteq T$ be an antichain which is maximal with respect to the property that $C_s \subseteq U_n$ for each $s \in A_n$. Since U_n is open dense in X, A_n will be a maximal antichain in T.

Let

$$V_n = \bigcup\{C_s : s \in A_n\}.$$

Then $V_n \subseteq U_n$ and V_n is open dense. (It is dense, because given any C_t there exists $s \in A_n$ and $r \in T$ with $t \leq r$ and $s \leq r$, hence $C_r \subseteq V_n \cap C_t$.)

Choose $\alpha < \omega_1$ so that for each $n \in \omega$ the (necessarily countable) antichain $A_n \subseteq T_{<\alpha}$. Let

$$U = \bigcup \{C_s : s \in T_\alpha\}.$$

Note that since T is normal U is an open dense set. Also

$$U \subseteq \bigcap_{n<\omega} V_n \subseteq \bigcap_{n<\omega} U_n.$$

($U \subseteq V_n$ because for any $b \in U$ if $b \in C_s$ for some $s \in T_\alpha$ there must be $t \in A_n$ comparable to it, since A_n is a maximal antichain, and since $A_n \subseteq T_{<\alpha}$, it must be that $t < s$ and so $b \in C_t \subseteq V_n$.

■

Lemma 4 *Suppose $f : X \to \mathbb{R}$ is a real valued Baire function. Then there exists $\alpha < \omega_1$ such that for each $s \in T_\alpha$ the function f is constant on C_s.*

Proof: Let $\mathcal{B}$ be a countable open basis for $\mathbb{R}$. For each $B \in \mathcal{B}$ the set $f^{-1}(B)$ has the property of Baire (open modulo meager). So there exists an open U_B such that

$$U_B \Delta f^{-1}(B) \text{ is meager.}$$

By the proof of Lemma 3 we may assume that

$$U_B = \bigcup \{C_s : s \in A_B\}$$

for some countable set $A_B \subseteq T$. By the proof of Lemma 3 there exists an $\alpha < \omega_1$ such that

- each $A_B \subseteq T_{<\alpha}$ and
- if U is the open dense set $\bigcup\{C_s : s \in T_\alpha\}$, then U is disjoint from $U_B \Delta f^{-1}(B)$ for each $B \in \mathcal{B}$.

But now, f is constant on each C_s for $s \in T_\alpha$. Otherwise, suppose that $f(b) \neq f(c)$ for some $b, c \in C_s$ for some $s \in T_\alpha$. Then suppose that $f(b) \in B$ and $f(c) \notin B$ for for some $B \in \mathcal{B}$. Because $b \in (f^{-1}(B) \cap U)$ and U is disjoint from $U_B \Delta f^{-1}(B)$, it must be that $b \in U_B$. Hence there exists $t \in T_{<\alpha}$ such

that $C_t \subseteq U_B$ and $b \in C_t$. Since $t < s$ it must be that $c \in C_t$ and so $c \in f^{-1}(B)$, which contradicts $f(c) \notin B$.

■

Steprans [S] shows that every continuous function on a Souslin tree takes on only countably many values.

Lemma 5 *Suppose $\{f_n : X \to \mathbb{R} : n \in \omega\}$ is a countable set of real valued Baire functions. Then there exists $\alpha < \omega_1$ such that for each $s \in T_\alpha$ and $n < \omega$ the function f_n is constant on C_s.*

Proof: Apply Lemma 4 countably many times and take the supremum of the α_n.

■

Finally, we prove the theorem. The idea of the proof is to use the argument of the atomic case, where the 'atoms' are supplied by Lemma 5. Since 'I-a.e. convergence' always implies 'convergence with respect to I', it is enough to see the converse. So let $f_n : X \to \mathbb{R}$ be Baire functions which converge to $f : X \to \mathbb{R}$ with respect to I, i.e. every subsequence has a subsequence which converges on a comeager set to f. By Lemma 5 there exists $\alpha < \omega_1$ such that for each $s \in T_\alpha$ and $n < \omega$ the function f_n is constant on C_s. It must be that for each fixed $s \in T_\alpha$ the constant values of f_n on C_s must converge on a comeager subset of C_s to a constant value. It follows that the sequence $f_n(x)$ converges to $f(x)$ on a comeager subset of the dense set $\{C_s : s \in T_\alpha\}$.

■

References

[CLO] K.Ciesielski, L.Larson, and K.Ostaszewski, **I-Density Continuous Functions**, Memoirs of the American Mathematical Society, 107(1994).

[PWW] W.Poreda, E.Wagner-Bojakoska, and W.Wilczyński, A category analogue of the density topology, Fundamenta Mathematicae, 125(1985), 167-173.

[S] J.Steprańs, Trees and continuous mappings into the real line, Topology and Its Applications, 12(1981), 181-185.

[ST] R.M.Solovay, S.Tennenbaum, Iterated Cohen extensions and Souslin's problem, Annals of Mathematics, 94(1971), 201-245.

[T] S.Todorčevič, Trees and linearly ordered sets, in **Handbook of Set Theoretic Topology**, North-Holland, (1984), 235-293.

[WW] E.Wagner, W.Wilczyński, Convergence of sequences of measurable functions, Acta Mathematica Academiae Scientiarum Hungaricae, 36(1980), 125-128.

Address: York University, Department of Mathematics, North York, Ontario M3J 1P3, Canada.

Permanent address: University of Wisconsin-Madison, Department of Mathematics, Van Vleck Hall, 480 Lincoln Drive, Madison, Wisconsin 53706-1388, USA).

e-mail: miller@math.wisc.edu

Nov 94